

Apalachicola (APA) NERR Nutrient Metadata
January – December 2014
Latest Update: February 1, 2018

I. Data Set and Research Descriptors

1) Principal investigators and contact persons

a) Reserve Contact

Rebecca Bernard, Research Coordinator
108 Island Drive
Eastpoint, FL 32328
850-670-7716
Rebecca.Bernard@dep.state.fl.us

Lauren Levi, Environmental Specialist II
108 Island Drive
Eastpoint, FL 32328
850-670-7710
lauren.levi@dep.state.fl.us

b) Laboratory Contact

Timothy W. Fitzpatrick
Chemistry Section
Florida Department of Environmental Protection
2600 Blair Stone Road
Tallahassee, FL 32399
850-245-8085

David D. Whiting
Biology Section
Florida Department of Environmental Protection
2600 Blair Stone Road
Tallahassee, FL 32399
850-245-8177

2) Research objectives

Previous studies have shown the importance of river flow and flushing rates on nutrients and primary productivity in the bay. Similar studies have determined nitrogen and phosphorus budgets for Apalachicola Bay as well as nutrient limitations related to seasonality and riverflow. There has been an ongoing controversy between the States of Florida, Georgia, and Alabama over the upstream diversion of water for 23 years. Approximately 88% of the drainage basin for the Apalachicola River and Bay is located in Georgia and Alabama and historical flows are being threatened by upstream development. A tri-state compact, between the states and approved by the US Congress, required negotiations between the states to develop a water allocation formula. The states were unable to come to an agreement and the compact expired. In late 2014 the US Supreme Court agreed to hear the case and legal proceedings are underway. This study is one of many looking at short-term variability, long-term change, and the relationship of other environmental factors to the productivity of the Apalachicola Bay system as

well as trying to separate natural from man-made variability. Data from this monitoring project has also been used by Florida DEP in support of Numeric Nutrient Criteria development.

a) Monthly Grab

Monthly grab samples are collected at 11 sites located across Apalachicola Bay to monitor spatial and temporal fluctuations in nutrient/chlorophyll *a* concentrations occurring in diverse sections of the bay. The stations have been chosen to help determine the influence of the river, local rainfall, adjacent habitats and anthropogenic impacts on these parameters. Sampling sites are located in the lower Apalachicola River, in the coastal area, offshore of the barrier islands, at the SWMP datalogger locations (primary SWMP stations), and throughout the bay. Seasonal, climatic, and anthropogenic factors all impact riverflow, which in turn affects nutrient/ chlorophyll *a* concentrations in the bay. Nutrient/chlorophyll *a* concentrations are also influenced by tidal action, wind direction and speed, and the hydrodynamics of the system.

b) Diel Sampling Program

Diel sampling is performed once a month in conjunction with grab sampling for nutrients/chlorophyll *a*. The East Bay Surface water quality datalogger site (apaesnut) is utilized each month for placement of the sampler so that temporal water quality data may be compared with the spatial nutrient/ chlorophyll *a* data collected at this site. Studies by the Reserve and others have shown the influence of tidal action and runoff on other physical parameters in the bay. Diel samples are collected over a 25 hour period thereby covering a lunar day of 24 hours 48 minutes.

3) Research methods

a) Monthly Grab Sampling Program

Monthly grab samples are collected at eleven stations (see Table 1) within and adjacent to Apalachicola Bay. All grab samples are collected on the same day. Due to the distance between the stations it is not always possible to collect all the samples several hours prior to low tide. Tidal condition, wind direction, speed, and cloud cover are recorded for each station at the time of sampling but are not included in this dataset and are available upon request. Climatic data from the ANERR weather station is available online at <http://cdmo.baruch.sc.edu/QueryPages/googlemap.cfm>. Sampling after heavy rains is avoided if at all possible. Water temperature, salinity, and dissolved oxygen are measured at surface and bottom for each station with a YSI 2030 handheld meter. pH is measured at surface and bottom for each station with a YSI 60 handheld meter. Surface measurements only are included in this dataset for temperature, salinity, pH, and dissolved oxygen (with the exception of the East Bay Bottom station). Bottom measurements for temperature, salinity, pH, and dissolved oxygen are available on request. Secchi data is also included in this dataset. Turbidity samples are collected at each site and are tested in the ANERR lab with a DRT-15CE Turbidimeter. All grab samples are analyzed at the Florida Department of Environmental Protection laboratory (FLDEP).

i) Grab sample collection:

A submersible pump and flexible clear plastic tubing is used to collect water from a depth of 0.5 meters at all stations not associated with a SWMP datalogger site. At the Cat Point and Dry Bar SWMP datalogger stations, water samples are collected at a depth of approximately 1.5 meters below the surface. This is the approximate depth of the probes of the data loggers deployed at these sites. At the East Bay datalogger station water samples are collected from surface (0.5 meters) and bottom (1.5 meters) depths, approximating the depths of the two dataloggers deployed at this site. Triplicate samples are collected every other month at one station, rotating through all station locations.

ii) Grab sample filtration and handling:

Water from the submersible pump is delivered directly into the appropriate sample bottles. A filter is attached to the end of the flexible tubing for collection of nutrient samples requiring filtration. Water filtered in this

manner is delivered directly to the appropriate sample bottles. Necessary preservatives are added prior to water sample according to appropriate EPA protocol for nutrient sampling. Whole water samples for chlorophyll *a* analysis are filtered at the FLDEP laboratory. All samples are placed on ice in the dark until delivery to the FLDEP laboratory. The submersible pump and tubing are flushed with ambient water prior to sample collection at each station. Filter holders are rinsed with DI prior to addition of a new filter at each station. A field blank is also run each month, using DI water for sample blank. The field blank is delivered using the pump, tubing and filter as described above. All grab samples are delivered to the FLDEP laboratory 24 to 36 hours after collection.

b) Diel Sampling Program

i) Diel sampling program:

Diel sampling is performed with an ISCO 3700 Portable Automated Sampler at the East Bay surface (apaesnut) station. The ISCO is deployed on a fixed platform located at the East Bay surface site. Generally the ISCO is deployed at the beginning of the grab sample collection trip and retrieved the following morning. In some months adverse weather conditions result in deployment of the ISCO sampler during a week other than the week of grab sample collection. The sampler is programmed to collect two 1000 ml water samples every 2.5 hours, over a 25-hour period at the same depth as the East Bay surface datalogger probes (1.7 m above the bottom sediment). This captures a complete 24 hr: 48min lunar-tidal cycle. The ISCO sampler is programmed to purge the suction line before and after each sample collection. The center of the ISCO sampler is filled with ice to aid in sample preservation. All samples are placed on ice upon retrieval of the ISCO sampler at the end of the sampling period. Nutrient sample filtration is performed at ANERR laboratory within one hour of retrieval from the ISCO sampler. Whole water samples for chlorophyll *a* analysis are filtered at the FLDEP laboratory. All diel samples are delivered to the FLDEP laboratory within 36 hours of the first sample collection time.

c) Equipment QAQC and maintenance – Grab and Diel Sampling Program:

The submersible pump, tubing, and field use filter holders are acid rinsed with 10% HCl and triple rinsed with DI water after each sampling trip. In lab use filtration funnels and receivers are acid washed with 10% HCl and triple rinsed with DI water after each sampling trip. Diel sample collection bottles used in the ISCO automated sampler are acid washed and triple with DI water after each sampling trip. The ISCO automated sampler tubing is acid washed and triple rinsed with DI water after each monthly sampling event. The overall condition of the pump and tubing is checked each month prior to deployment, tubing is replaced as needed. New unused sample bottles are supplied by FLDEP laboratory for each sampling event. The YSI 2030, pH meter, and Turbidimeter are calibrated each day of use.

4) Site location and character

The Apalachicola Drainage Basin encompasses over 50,700 square kilometers and includes parts of three states (Alabama, Georgia, and Florida). The Apalachicola River is the largest in Florida in terms of flow. The amount of river discharge has been shown to be highly significant to the ecology of the estuary, which acts as a buffer between the Gulf of Mexico and fresh water input from upland areas. The nutrient rich plume of "green water" moving out of Apalachicola Bay is also important to the productivity of the northeastern Gulf of Mexico. The Apalachicola National Estuarine Research Reserve is located in the northwestern part of Florida, generally called the panhandle. It is located adjacent to the City of Apalachicola, and encompasses most of the Apalachicola Bay system, including 52 miles of the lower Apalachicola River. Passes, both natural and manmade, connect Apalachicola Bay to the northeastern Gulf of Mexico.

Monthly grab samples are collected at all nutrient monitoring stations. A map of station locations is given in Figure 1.

a) East Bay datalogger and nutrient station

East Bay is separated from Apalachicola Bay by two bridges and a causeway and is located to the north of the bay proper. East Bay is 8.2 km long, has an average Depth of approximately 1.0 m MHW, and an average width of 1.8 km. The tides in East Bay are mixed and range from 0.3 m to 1.0 m (average 0.5 m). The datalogger and nutrient sampling site is located in the upper reaches of East Bay. The piling location for the two East Bay dataloggers (ES and EB) is latitude 29°47.15' N and longitude 84°52.52' W (29.7858 N, 84.8752 W). At the sampling site, the Depth is 2.2 m MHW and the width of the bay is 1 km. The tides in the system are mixed, meaning the number of tides can range from one to five tides during a 24 hour period and are not evenly distributed throughout the day. At the East Bay bottom site the meter probes are 0.3 m above the bottom sediment. Salinity ranges from 0 to 30 ppt and the long-term (1993-2013) average salinity is 10.6 ppt. At the East Bay surface site the meter probes are 1.7 m above the bottom sediment and salinity ranges from 0 ppt to 30 ppt with a long term (1995-2013) average salinity of 9.6 ppt. The freshwater input is very tannic and usually dark colored. Flows vary with local rainfall and are not quantified due to the diverse sources of the runoff. The bottom habitat at this bay site is soft sediment, primarily silt and clay, with no vegetation present. The dominant marsh vegetation near the sampling site (approximately 300 meters away) is needlerush grass (*Juncus roemerianus*) and swamp sawgrass (*Cladium jamaicense*) and smooth cordgrass (*Spartina alterniflora*). The dominant upland vegetation is primarily pineland forests which includes slash pine (*Pinus elliotii*), saw palmetto (*Serenoa repens*), and sand pine (*Pinus clausa*). Upland land use near the sampling site includes conservation and silviculture uses with some single family residential in the lower East Bay area. The sampling site is influenced by local runoff from Tate's Hell Swamp, the East Bay marshes, and distributary flow, some of which comes from the Apalachicola River via the East River. Tate's Hell Swamp was ditched, diked, and altered in the late 1960's and early 1970's by timber companies. These changes shortened the drainage period and allowed increased runoff with a concomitant decrease in pH and increase in color, which had a drastic effect on the biological communities in East Bay. Restoration of Tate's Hell Swamp began in 1995 to reduce non-point source runoff and restore historic sheet flow in the area.

b) Cat Point datalogger and nutrient station

The Cat Point datalogger and nutrient sampling site is located in St. George Sound, approximately 400 meters east of the St. George Island Bridge. The piling location is latitude 29°42.12' N and longitude 84°52.81' W (29.7021 N, 84.8802 W). The tides at Cat Point are mixed and range from 0.3 m to 1.0 m (average 0.5 m). At the sampling site, the depth is 2.5 m MHW. (The site was moved approximately 600 meters south in October 1997) and the width of the bay is 6.4 kilometers. At the Cat Point site the meter probes are 0.3 meters above the bottom sediment. This is also the depth where nutrients are collected monthly. Salinity ranges from 0 to 32 ppt with a long-term (1992-2013) average salinity of 21.4 ppt. Flows vary with local rainfall and are not quantified due to the diverse sources of the runoff. The bottom type is oyster bar with no vegetation present except algae growing on the oysters in the summer. The dominant upland vegetation is primarily pineland forests, which include slash pine (*Pinus elliotii*), saw palmetto (*Serenoa repens*), and sand pine (*Pinus clausa*). Upland land use near the sampling site includes single family residential and commercial use in the Eastpoint area. The sampling site is influenced by local runoff from Tate's Hell Swamp and flow from the Apalachicola River. High salinity water comes mainly from the east, through East Pass at the eastern end of St. George Island.

c) Dry Bar datalogger and nutrient station

The Dry Bar datalogger and nutrient sampling site is located near St. Vincent Sound, in the western part of the Apalachicola Bay system, approximately 0.8 kilometer east of St. Vincent Island. The piling location is latitude 29°40.48' N and longitude 85°03.50' W (29.6747 N, 85.0583 W). At the sampling site, the depth is 2 meters and the width of the bay is 11.2 kilometers. At the Dry Bar site the datalogger probes are located 0.3 meters above the bottom sediment. This is also the depth where nutrients are collected monthly. The tides are mixed and range

from 0.3 to 1.0 meters. Salinity ranges from 0 to 34 ppt with a long-term (1992-2013) average salinity of 20.9 ppt. The bottom type is oyster bar with no vegetation present, except algae that grows on the oysters during the summer months. The dominant upland vegetation includes slash pine (*Pinus clausa*) flatwoods with various combinations of gallberry (*Ilex glabra*), smooth cordgrass (*Spartina alterniflora*), fetterbush (*Leucothoe racemosa*), cabbage palm (*Sabal palmetto*), saw palmetto (*Serenoa repens*), magnolia (*Magnolia grandiflora*), and grasses. Upland use near the sampling site includes state owned and managed Cape St. George Island, St. Vincent National Wildlife Refuge, as well as single family residential and commercial use in the Apalachicola area. The sampling site is influenced by the flow of the Apalachicola River and high salinity water coming through West Pass and Sikes Cut.

d) Secondary SWMP stations

Detailed information for the additional 7 nutrient (secondary SWMP) stations, not associated with the required sampling at the datalogger sites, as well as the datalogger sites, is included in Table 1.

West Pass

29.6379 N, 85.089 W

Salinity ave = 22.5 ppt, range = 1.8-36.0 ppt

This site is located in the pass between two uninhabited barrier islands, the state owned and managed Cape St. George Island and St. Vincent National Wildlife Refuge. The sampling site is influenced by the flow of the Apalachicola River and high salinity water coming through West Pass.

Pilots Cove

29.6079 N, 85.0196 W

Salinity ave = 22.9 ppt, range = 1.3-35.5 ppt

This site is located near state owned and managed Cape St. George Island, an uninhabited barrier island. The sampling site is influenced by the flow of the Apalachicola River and high salinity water coming through West Pass.

Mid Bay

29.6677 N, 84.994 W

Salinity ave = 16.3 ppt, range = 0.2-35.2 ppt

This sampling site is located in central Apalachicola Bay. The site is roughly equidistant from state owned and managed Cape St. George Island (four miles distant), St. Vincent National Wildlife Refuge (six miles distant), and single family residential and commercial use in the Apalachicola area (four miles distant). This site is approximately 1.5 miles from the intracoastal waterway channel. The sampling site is influenced by the flow of the Apalachicola River and high salinity water coming through Sikes Cut and West Pass.

East Bay Bridge

29.7308 N, 84.9452 W

Salinity ave = 7.9 ppt, range = 0-30.7 ppt

This site is located near the western section of the US Highway 98 bridge connecting Apalachicola and Eastpoint. The bridge also serves as the boundary line between East Bay and Apalachicola Bay. Nearby upland areas consist of residential and commercial use in Apalachicola and Eastpoint areas. The sampling site is influenced by flows from the Apalachicola River and tributaries including the Little St Marks river, St. Marks river, and East river.

Sikes Cut offshore

29.6067 N, 84.9467 W

Salinity ave = 31.9 ppt, range 21.7-35.8 ppt

This site is selected to characterize true marine water, and is located south of Sikes Cut in the Gulf of Mexico. The site is near the eastern portion of state owned and managed Cape St. George Island and near the western end

of St. George Island in an area consisting of single family and vacation homes. Sikes Cut allows tidal exchange of high salinity water from the Gulf of Mexico and lower salinity water from Apalachicola Bay. Sikes Cut is an important pass utilized by commercial and recreational vessels.

Nicks Hole

29.6504 N, 84.9289 W

Salinity ave = 19.0, range = 0.5-35.4 ppt

This site is near single family and vacation home use on St George Island. A small airport utilized by private aircraft is also located near Nicks Hole. The site is tidally influenced by high salinity water from Sikes Cut and by flows from the Apalachicola River.

River

29.7791 N, 85.0434 W

Salinity ave = 0.1, range = 0-0.1 ppt


This site is selected to characterize fresh water in the Apalachicola River. The site is located in the central channel of the river approximately 6 miles north and upstream of the river mouth and the residential and commercial areas of Apalachicola. Adjacent areas are state owned and managed forested floodplain. The site is influenced by Apalachicola River flow.

Table 1. Nutrient and chlorophyll *a* sampling sites for the Apalachicola NERR SWMP.

Station code	Station name	Latitude	Longitude	Tidal range average (meters)	Salinity range	Water depth average (meters)	Bottom habitat	Datalogger station name	Sample depth (meters)
apawpnut	West Pass	29 38.279	85 5.341	0.7	euryhaline	5.0	sand		0.5
apadbnut	Dry Bar	29 40.482	85 3.502	0.7	euryhaline	1.7	oyster bar	apadb	1.5
apapcnut	Pilot's Cove	29 36.473	85 1.173	0.7	euryhaline	1.8	patchy seagrass		0.5
apambnut	Mid Bay	29 40.061	84 59.641	0.7	euryhaline	2.2	sandy silt		0.5
apaegnut	East Bay Bridge	29 43.848	84 56.711	0.7	euryhaline	1.6	silty clay		0.5
apaesnut	East Bay Surface	29 47.147	84 52.512	0.7	euryhaline	1.7	clayey sand	apaes	0.5
apaebnut	East Bay Bottom	29 47.147	84 52.512	0.7	euryhaline	1.7	clayey sand	apaeb	1.5
apascnut	Sikes Cut Offshore	29 36.401	84 56.799	0.7	marine	>5.0	sand		0.5
apanhnut	Nick's Hole	29 39.022	84 55.732	0.7	euryhaline	1.0	patchy seagrass		0.5
apacpnut	Cat Point	29 42.128	84 52.811	0.7	euryhaline	1.8	oyster bar	apacp	1.5
aparvnut	River	29 46.743	85 2.606	0.7	oligohaline	3-4	sandy silt		0.5

Note: Diel samples are collected 2.5 hours apart at the East Bay Surface datalogger site, APAESNUT, with the ISCO automated water sampler. No duplicate diel samples are taken, however there is some overlap with monthly grabs collected at the East Bay Surface station and deployment of the ISCO sampler.

Figure 1. Station locations.



5) Code variable definitions

Station code names:

apacpnut = Apalachicola Reserve nutrient data for Cat Point
 apadbnut = Apalachicola Reserve nutrient data for Dry Bar
 apaebnut = Apalachicola Reserve nutrient data for East Bay Bottom
 apaegnut = Apalachicola Reserve nutrient data for East Bay Bridge
 apaesnut = Apalachicola Reserve nutrient data for East Bay Surface
 apambnut = Apalachicola Reserve nutrient data for Mid Bay
 apanhnut = Apalachicola Reserve nutrient data for Nicks Hole
 apapcnut = Apalachicola Reserve nutrient data for Pilots Cove

aparvnut = Apalachicola Reserve nutrient data for River
 apascnut = Apalachicola Reserve nutrient data for Sikes Cut
 apawpnut = Apalachicola Reserve nutrient data for West Pass

Monitoring Programs:

Monthly grab samples = 1

Diel grab sampling = 2

6) Data collection period

Nutrient monitoring began in April 2002 at all stations listed. Sampling has been performed monthly at all stations, unless otherwise noted. This table lists collection times for all nutrient and chlorophyll *a* samples in 2014. The below Start and End time reflect the times that the first and last diel samples were collected for each monthly diel sampling event. Grab sample end time is not recorded in the field. Generally grab sample collection, filtering, and icing are completed within 10 minutes or less depending upon field conditions at the time of sampling. Time is coded based on a 2400 hour clock and is referenced to Eastern Standard Time (EST), without Daylight Savings Time adjustments.

a) Samples date/times for Monitoring Program 1 (Grab Samples)

Site	Date/Time	Site	Date/Time
apacpnut	01/13/2014 11:11	apadbnut	01/13/2014 09:10
apacpnut	02/10/2014 10:05	apadbnut	02/10/2014 12:39
apacpnut	03/04/2014 10:14	apadbnut	03/04/2014 12:16
apacpnut	04/01/2014 08:58	apadbnut	04/01/2014 10:33
apacpnut	05/06/2014 10:15	apadbnut	05/06/2014 11:50
apacpnut	06/10/2014 10:16	apadbnut	06/10/2014 07:36
apacpnut	07/01/2014 12:56	apadbnut	07/01/2014 09:13 - 09:21
apacpnut	08/06/2014 09:10	apadbnut	08/06/2014 11:22
apacpnut	09/03/2014 09:30	apadbnut	09/03/2014 11:39
apacpnut	10/07/2014 09:10	apadbnut	10/07/2014 11:42
apacpnut	10/28/2014 08:55	apadbnut	10/28/2014 10:58
apacpnut	12/03/2014 09:00	apadbnut	12/03/2014 10:25

Site	Date/Time	Site	Date/Time
apaebnut	01/13/2014 11:39	apaegnut	01/13/2014 12:10
apaebnut	02/10/2014 10:40	apaegnut	02/10/2014 09:46
apaebnut	03/04/2014 09:01 - 09:04	apaegnut	03/04/2014 09:57
apaebnut	04/01/2014 08:05	apaegnut	04/01/2014 08:40
apaebnut	05/06/2014 09:17 – 09:34	apaegnut	05/06/2014 09:58
apaebnut	06/10/2014 09:34	apaegnut	06/10/2014 09:52
apaebnut	07/01/2014 08:06	apaegnut	07/01/2014 08:37
apaebnut	08/06/2014 08:17	apaegnut	08/06/2014 08:50
apaebnut	09/03/2014 08:02	apaegnut	09/03/2014 09:15
apaebnut	10/07/2014 08:20	apaegnut	10/07/2014 08:53
apaebnut	10/28/2014 08:07	apaegnut	10/28/2014 08:42
apaebnut	12/02/2014 10:21	apaegnut	12/02/2014 10:33

Site	Date/Time	Site	Date/Time
apaesnut	01/13/2014 11:37	apambnut	01/13/2014 10:09
apaesnut	02/10/2014 10:38	apambnut	02/10/2014 12:59
apaesnut	03/04/2014 08:58	apambnut	03/04/2014 12:30
apaesnut	04/01/2014 08:02	apambnut	04/01/2014 10:11
apaesnut	05/06/2014 09:15	apambnut	05/06/2014 12:15
apaesnut	06/10/2014 09:23	apambnut	06/10/2014 07:17
apaesnut	07/01/2014 08:03	apambnut	07/01/2014 08:52
apaesnut	08/06/2014 08:14	apambnut	08/06/2014 09:30
apaesnut	09/03/2014 08:00	apambnut	09/03/2014 11:56
apaesnut	10/07/2014 08:15	apambnut	10/07/2014 12:10
apaesnut	10/28/2014 08:00	apambnut	10/28/2014 11:31
apaesnut	12/02/2014 10:14	apambnut	12/03/2014 10:05
Site	Date/Time	Site	Date/Time
apanhnut	01/13/2014 10:42	apapcnut	01/13/2014 09:55
apanhnut	02/10/2014 11:28	apapcnut	02/10/2014 12:07
apanhnut	03/04/2014 10:33	apapcnut	03/04/2014 11:41
apanhnut	04/01/2014 09:18	apapcnut	04/01/2014 11:07
apanhnut	05/06/2014 10:38	apapcnut	05/06/2014 11:15
apanhnut	06/10/2014 08:47	apapcnut	06/10/2014 08:11
apanhnut	07/01/2014 12:35	apapcnut	07/01/2014 11:54
apanhnut	08/06/2014 09:45	apapcnut	08/06/2014 10:40
apanhnut	09/03/2014 09:51	apapcnut	09/03/2014 10:53 – 10:58
apanhnut	10/07/2014 09:32	apapcnut	10/07/2014 10:33
apanhnut	10/28/2014 09:15	apapcnut	10/28/2014 09:58
Apanhnut	*missing sample	apapcnut	*missing sample
Site	Date/Time	Site	Date/Time
aparvnut	01/13/2014 12:38 – 12:45	apascnut	*missing sample
aparvnut	02/10/2014 09:17	apascnut	02/10/2014 11:47
aparvnut	03/04/2014 13:00	apascnut	03/04/2014 11:11
aparvnut	04/01/2014 11:42	apascnut	04/01/2014 09:47
aparvnut	05/06/2014 12:41	apascnut	05/06/2014 10:56
aparvnut	06/10/2014 12:00	apascnut	06/10/2014 08:32
aparvnut	07/01/2014 07:15	apascnut	07/01/2014 12:19
aparvnut	08/06/2014 12:10	apascnut	08/06/2014 10:10
aparvnut	09/03/2014 12:35	apascnut	09/03/2014 10:10
aparvnut	10/07/2014 12:40	apascnut	10/07/2014 10:08
aparvnut	10/28/2014 12:04	apascnut	10/28/2014 09:34
aparvnut	12/02/2014 09:02	apascnut	12/03/2014 09:42

Site	Date/Time
apawpnut	01/13/2014 09:38
apawpnut	02/10/2014 12:20
apawpnut	03/04/2014 11:59
apawpnut	04/01/2014 10:52
apawpnut	05/06/2014 11:30
apawpnut	06/10/2014 07:58
apawpnut	07/01/2014 11:37
apawpnut	08/06/2014 10:58
apawpnut	09/03/2014 11:13
apawpnut	10/07/2014 11:11
apawpnut	10/28/2014 10:33 – 10:46
apawpnut	*missing sample

b) Start and End Date/Time for Monitoring Program 2 (Diel Sampling)

Site	Start Date/Time	End Date/Time
apaesnut	01/09/2014 10:45	01/10/2014 11:45
apaesnut	02/10/2014 11:00	02/11/2014 12:00
apaesnut	03/04/2014 09:45	03/05/2014 10:45
apaesnut	04/01/2014 08:15	04/02/2014 09:15
apaesnut	05/06/2014 10:00	05/07/2014 11:00
apaesnut	06/10/2014 09:30	06/11/2014 11:30
apaesnut	07/01/2014 08:30	07/02/2014 09:30
apaesnut	08/06/2014 08:15	08/07/2014 09:15
apaesnut	09/03/2014 08:30	09/04/2014 09:30
apaesnut	10/07/2014 09:00	10/08/2014 10:00
apaesnut	10/28/2014 08:30	10/29/2014 09:30
apaesnut	12/02/2014 10:15	12/03/2014 11:15

7) Associated researchers and projects

The Reserve conducts long-term water quality monitoring and maintains a weather station as part of the NERRS SWMP. SWMP water quality and weather data may be accessed via www.nerrsdata.org.

Other ongoing projects or data that relate to the nutrient monitoring project include:

Apalachicola Bay Oyster Situation Report TP200, UF/IFAS, Sea Grant Florida, April 24, 2013.

Apalachicola River Discharge, U.S. Geological Survey, <http://waterdata.usgs.gov/nwis/>

Garland, H., Christopher, M., Cox, N., Yuan, S.,
Apalachicola National Estuarine Research Reserve
Oyster reef research in Apalachicola Bay provided for the ACF lawsuit.

Byars, Natalie, Florida State University

How does climatic- and human-induced variability in river flow affect the spatial-temporal distribution of phytoplankton and their subsequent availability to oysters in Apalachicola Bay, Florida?

Caffrey, Jane, University of West Florida

Effect of Diurnal and Weekly Water Column Hypoxic Events on Nitrification and Nitrogen Transformations in Estuarine Sediments

Hagen, S., DeLorme, D., Walters, L., Wang, D., Weishampel, J., Yeh, G., Huang, W., Slinn, D., Morris, J.
Ecological Effects of Sea Level Rise

“Gauging the effects of the BP Oil Spill on diatoms, calcareous nanoplankton, and related protists at or near the base of the food chain in the NE Gulf of Mexico”, funded to principal Investigators, Drs Sherwood W. Wise, Jr. and Akshitnala K. S. K. Prasad.

Garwood, J., Harper, J., Lamb, M., Jones, D., Levi, L., Bourque, E.

Apalachicola National Estuarine Research Reserve.

Distribution and density of fishes and benthic invertebrates in Apalachicola Bay.

Harper, J., Wren, K., Jones, D., Garwood, J., Canedo, J., Snyder, C., Levi, L./ NERRS Sentinel Sites Program for Understanding Climate Change Impacts on Estuaries

Jones, D., Bourque, E., Lamb, M., Harper, J., Levi, L., Garwood, J.

Apalachicola National Estuarine Research Reserve

System Wide Monitoring Program

Long-Term Water Quality Monitoring

Levi, L., Harper, J., Garwood, J.

Apalachicola National Estuarine Research Reserve

System Wide Monitoring Program

Long-Term Meteorological Monitoring

Site-Specific Information in Support of Establishing Numeric Nutrient Criteria in Apalachicola Bay, Nutrient Criteria Technical Support Document. Division of Assessment and Restoration, Florida Department of Environmental Protection, July, 2013.

Tucker, Kim, Florida Agricultural and Mechanical University, Department of Civil and Environmental Engineering, Master’s Thesis, Effects of river flow and rainfall on chlorophyll a in Apalachicola River. 2011.

Viveros, Paula, NOAA Graduate Research Fellowship, University of Florida

Phytoplankton composition and abundance in relation to salinity, nutrient and light gradients in the Apalachicola National Estuarine Research Reserve (ANERR)

Wang, H., W. Huang, M. Harwell, L. Edmiston, E. Johnson, P. Hsieh, K. Milla, J. Christensen,

J. Stewart, X. Liu. 2008. Modeling oyster growth rate by coupling oyster population and hydrodynamic models for Apalachicola Bay, Florida, USA. Ecological Modeling 211:77-89.

8) Distribution

NOAA retains the right to analyze, synthesize and publish summaries of the NERRS System-wide Monitoring Program data. The NERRS retains the right to be fully credited for having collected and process the data. Following academic courtesy standards, the NERR site where the data were collected should be contacted and fully acknowledged in any subsequent publications in which any part of the data are used. The data set enclosed within this package/transmission is only as good as the quality assurance and quality control procedures outlined by the enclosed metadata reporting statement. The user bears all responsibility for its subsequent use/misuse in any further analyses or comparisons. The Federal government does not assume liability to the Recipient or third persons, nor will the Federal government reimburse or indemnify the Recipient for its liability due to any losses resulting in any way from the use of this data.

Requested citation format:

NOAA National Estuarine Research Reserve System (NERRS). System-wide Monitoring Program. Data accessed from the NOAA NERRS Centralized Data Management Office website: www.nerrsdata.org; *accessed* 12 October 2012.

NERR nutrient data and metadata can be obtained from the Research Coordinator at the individual NERR site (please see Principal investigators and contact persons), from the Data Manager at the Centralized Data Management Office (please see personnel directory under the general information link on the CDMO home page) and online at the CDMO home page www.nerrsdata.org. Data are available in comma separated version format

II. Physical Structure Descriptors

9) Entry verification

A hard copy of the FLDEP Central Laboratory Sample Submittal Form accompanies the samples from ANERR to the FLDEP laboratory. FLDEP laboratory results and final reports are transmitted to ANERR by electronic download via the FLDEP Laboratory Information Management System (LIMS). Analysis data is exported from LIMS in Excel format and is then formatted by Lauren Levi, ANERR staff, according NERR CDMO SOPs. Original FLDEP LIMS downloaded data files, formatted data files and field logs are all compared with each other to ensure accuracy.

Nutrient data are entered into a Microsoft Excel worksheet and processed using the NutrientQAQC Excel macro. The NutrientQAQC macro sets up the data worksheet, metadata worksheets, and MDL worksheet; adds chosen parameters and facilitates data entry; allows the user to set the number of significant figures to be reported for each parameter and rounds using banker's rounding rules; allows the user to input MDL values and then automatically flags/codes measured values below MDL and inserts the MDL; calculates parameters chosen by the user and automatically flags/codes for component values below MDL, negative calculated values, and missing data; allows the user to apply QAQC flags and codes to the data; produces summary statistics; graphs selected parameters for review; and exports the resulting data file to the CDMO for tertiary QAQC and assimilation into the CDMO's authoritative online database.

Lauren Levi was responsible for these tasks.

10) Parameter Titles and Variable Names by Data Category

Required NOAA/NERRS System-wide Monitoring Program nutrient parameters are denoted by an asterisk ”*”.

Data Category	Parameter	Variable Name	Units of Measure	Collection Period
Phosphorus:	*Orthophosphate, filtered	PO4F	mg/L as P	Jan-Dec 2014
	Total Phosphorus	TP	mg/L as P	Jan-Dec 2014
Nitrogen:	*Nitrite + Nitrate, filtered	NO23F	mg/L as N	Jan-Dec 2014
	*Ammonium, filtered	NH4F	mg/L as N	Jan-Dec 2014
	Dissolved Inorganic Nitrogen	DIN	mg/L as N	Jan-Dec 2014
	Total Kjeldahl Nitrogen whole	TKN	mg/L as N	Jan-Dec 2014
	Total Nitrogen	TN	mg/L as N	Jan-Dec 2014
Plant Pigments:	*Chlorophyll <i>a</i>	CHLA_N	µg/ L	Jan-Dec 2014
	Uncorrected Chlorophyll <i>a</i>	UncCHLA_N	µg/L	Jan-Dec 2014
	Phaeophytin	PHEA	µg/ L	Jan-Dec 2014
Other Laboratory Parameters:	Total Suspended Solids	TSS	mg/L	Jan-Dec 2014
Field Parameters:	Water temperature	WTEM_N	°C	Jan-Dec 2014
	Salinity	SALT_N	ppt	Jan-Dec 2014
	Dissolved oxygen	DO_N	mg/L	Jan-Dec 2014
	% Saturated dissolved oxygen	DO_S_N	%	Jan-Dec 2014
	pH	PH_N	SU	Jan-Dec 2014
	Turbidity	TURB_N	NTU	Jan-Dec 2014
	Secchi Disk Depth	SECCHI	meters	Jan-Dec 2014

Notes:

1. Time is coded based on a 2400 hour clock and is referenced to Standard Time.
2. Reserves have the option of measuring either NO₂ and NO₃ or they may substitute NO₂ for individual analyses if they can show that NO₂ is a minor component relative to NO₃. ANERR has shown NO₂ to be a minor component of NO₂3.

11) Measured and Calculated Laboratory Parameters

a) Parameters Measured Directly

Nitrogen species:	NO23F, NH4F, TKN
Phosphorus species:	PO4F, TP
Other:	UncCHLA_N, CHLA_N, PHEA, TSS

b) Calculated Parameters

DIN:	NO23F+NH4F
TN:	NO23F + TKN

12) Limits of Detection

All information in this section is provided by FLDEP laboratory.

a) FLDEP laboratory MDL determination:

Minimum detection limits (MDLs) are set such that the risk of reporting a false positive is less than 1%. MDLs are determined using the method specified in the Federal Register, 40 CFR Part 136 Appendix B, using LCS (laboratory control samples) prepared near the estimated detection limit as surrogates to estimate methodological noise for censored methods (e.g., chromatographic methods which censor analytical noise) or, for uncensored methods, using actual method blanks to directly measure methodological noise. Where the possibility exists for significant systematic bias from sample preparation and handling or from the analytical determinative step (typically inorganic analyses), bias is taken into account when calculating detection limits. Published MDLs may be set higher than experimentally determined MDLs to (1) avoid observed positive interferences from matrix effects or common reagent contaminants or (2) for reporting convenience (i.e., to group common compounds with similar but slightly different experimentally determined MDLs). MDLs are determined in a suitable analyte-free matrix when possible. For certain analytes and matrices, no suitable, analyte-free matrix may be available. In those cases, MDLs are determined in the absence of any matrix, but in the presence of all preparatory reagents carried through the full preparatory and determinative steps. Level of detection (LOD) verification procedures may be found in SOP LB-031, Limit of Detection Verification. (From page 39 of FLDEP Laboratory Quality Manual 2014 http://www.FLDEP.state.fl.us/labs/docs/lab_qualitymanual14.pdf)

Table 3. Method Detection Limits for FLDEP laboratory. Note that the FLDEP laboratory reports the MDL value for all analytical results that fall below MDL.

Parameter	Start Date	End Date	MDL	UNITS
NH4F	01/01/14	12/31/14	0.002	mg/L as N
NO23F	01/01/14	12/31/14	0.004	mg/L as N
PO4F	01/01/14	12/31/14	0.004	mg/L as P
TP	01/01/14	12/31/14	0.005	mg/L as P
TKN	01/01/14	12/31/14	0.08	mg/L as N
TSS	01/01/14	12/31/14	2.0	mg/L

FLDEP MDLs for the chlorophyll suite of components may change by station and month based on the need to dilute samples during processing. The base MDL listed in the FLDEP SOP is based on the maximum filtration volume and minimum extract volume and will therefore be the lowest MDL.

Base MDL values for ANERR 2014 plant pigment parameters:

Parameter	Start Date	End Date	MDL	Units
CHLA_N	01/01/14	12/31/14	0.55	ug/L
UncCHLA_N	01/01/14	12/31/14	0.40	ug/L
PHEA	01/01/14	12/31/14	0.40	ug/L

The sample MDL is calculated based on the number of times a sample must be diluted. For example, if a CHL_A sample must be diluted to twice its volume, the base MDL of 0.55 ug/L is multiplied by a dilution factor of two (0.55 ug/L x 2) thus resulting in an MDL of 1.10 ug/L. For samples that fall

below the MDL and their MDL is greater than the base MDL, individual sample MDLs are listed in the table below. These data have been flagged and coded as -4 SBL in the dataset.

2014 MDLs for Chlorophyll *a* (CHLA_N), Uncorrected Chlorophyll *a* (UncCHLA_N), and Phaeophytin (PHEA), as reported by FLDEP laboratory when values differ from base MDL values:

Component	DateTimeStamp	Site	MDL	Units
CHLA_N	04/01/2014 10:45	apaesnut	0.85	ug/L
CHLA_N	05/06/2014 09:17	apaebnut	0.92	ug/L
CHLA_N	05/06/2014 09:32	apaebnut	0.85	ug/L
CHLA_N	05/06/2014 09:34	apaebnut	0.79	ug/L
CHLA_N	05/06/2014 10:00	apaesnut	0.79	ug/L
CHLA_N	05/06/2014 15:00	apaesnut	0.79	ug/L
CHLA_N	05/06/2014 17:30	apaesnut	1.1	ug/L
CHLA_N	05/06/2014 20:00	apaesnut	0.79	ug/L
CHLA_N	05/06/2014 22:30	apaesnut	1.1	ug/L
CHLA_N	05/07/2014 01:00	apaesnut	0.79	ug/L
CHLA_N	05/07/2014 03:30	apaesnut	0.92	ug/L
CHLA_N	05/07/2014 06:00	apaesnut	0.92	ug/L
CHLA_N	05/07/2014 08:30	apaesnut	1.0	ug/L
CHLA_N	05/07/2014 11:00	apaesnut	1.2	ug/L
PHEA	01/13/2014 12:38	aparvnut	0.44	ug/L
PHEA	02/10/2014 11:00	apaesnut	0.80	ug/L
PHEA	02/10/2014 18:30	apaesnut	0.80	ug/L
PHEA	05/06/2014 09:17	apaebnut	0.67	ug/L
PHEA	05/06/2014 09:32	apaebnut	0.62	ug/L
PHEA	05/06/2014 12:30	apaesnut	0.67	ug/L
PHEA	05/06/2014 17:30	apaesnut	0.83	ug/L
PHEA	05/06/2014 22:30	apaesnut	0.80	ug/L
PHEA	05/07/2014 01:00	apaesnut	0.57	ug/L
PHEA	05/07/2014 03:30	apaesnut	0.67	ug/L
PHEA	05/07/2014 06:00	apaesnut	0.67	ug/L
PHEA	05/07/2014 08:30	apaesnut	0.73	ug/L
PHEA	07/01/2014 11:37	apawpnut	0.49	ug/L
PHEA	07/01/2014 08:37	apaegnut	0.75	ug/L
PHEA	07/01/2014 12:35	apanhnut	0.55	ug/L

13) FLDEP Laboratory Methods

a) Parameter: PO4

- 1) Method Reference: The described procedure is based on EPA Method 365.1, Rev. 2.0 (1993) and the Bran+Lubbe method G-146-95 Rev. 3.
- 2) Method Description: Orthophosphate reacts with molybdenum (VI) and antimony (III) in an acid medium to form an antimony-phospho-molybdate complex. The complex is reduced with ascorbic acid to form a blue complex that absorbs at 880 nm.
- 3) Preservation Method: Samples are filtered in the field, placed on ice in the dark and analyzed within 48 hours.

b) Parameter: TP

- 1) Method Reference: This SOP is based on EPA Method 365.1, Rev. 2.0 (1993) and SEAL Analytical AQ2 Method: EPA-119-A Rev. 5.
- 2) Method Description: Prior to analysis the samples are acid-persulfate digested according to the DEP SOP NU-049. This process converts inorganic and organic forms of phosphorus to ortho-phosphate. Ortho-phosphate reacts with molybdenum and antimony in an acidic medium to form a phosphoantimony/molybdenum complex, which is reduced with ascorbic acid. The AQ2 Discrete Analyzer is used to measure the absorbance of the complex at 880 nm.
- 3) Preservation Method: Whole water is acidified in the field to pH <2, placed on ice in the dark and analyzed within 28 days.

c) Parameter: NH4

- 1) Method Reference: This SOP is based upon EPA Method 350.1, Rev. 2.0 (1993) and OI Analytical Method 3271152 utilizing gas diffusion.
- 2) Method Description: The sample pH is raised to a pH of >11. The ammonia molecules generated at this pH pass through a gas diffusion membrane and are absorbed into an alkaline hypochlorite solution to form chloramine. The chloramine reacts with salicylate to form indophenol blue in an amount that is proportional to the ammonia concentration. Sodium nitroferrocyanide intensifies the blue color. The absorbance is measured at 660 nm.
- 3) Preservation Method: Samples are filtered in the field, acidified to pH <2, placed on ice in the dark and analyzed within 28 days.

d) Parameter: NO23

- 1) Method Reference: This method is based on EPA method 353.2, Rev 2.0 (1993) and Lachat method 10-107-04-1-C.
- 2) Method Description: A filtered sample is passed through a column containing granular copper-cadmium, which reduces nitrate to nitrite. The nitrite originally present plus the reduced nitrate can then be determined by colorimetry. The nitrite is diazotized with sulfanilamide and coupled with N-(1-naphthyl) ethylenediamine dihydrochloride to form a highly colored azo dye, which is measured at a wavelength of 520 nm.
- 3) Preservation Method: Samples are filtered in the field, acidified to pH <2, placed on ice in the dark and analyzed within 28 days.

e) Parameter: TKN

- 1) Method Reference: This SOP is based on EPA method 351.2, Rev. 2.0 (1993) and Seal AQ2 method EPA-111-A Rev. 4.
- 2) Method Description: Prior to analysis, digestion converts free ammonia and organic nitrogen compounds to ammonium sulfate (DEP SOP NU-091). Ammonium reacts with salicylate and hypochlorite in a buffered, alkaline solution in the presence of sodium nitroferrocyanide (pH = 12.4-12.7) to form the salicylic acid analog of indophenol blue. The blue-green color produced is measured at 660 nm.

- 3) Preservation Method: Whole water is acidified in the field to pH <2, placed on ice in the dark and analyzed within 28 days.

f) Parameter: CHLA_N and UncCHLA_N and PHEA

- 1) Method Reference: This method is based on Standard Methods 10200H and EPA Method 446.0.
- 2) Method Description: This method is used to determine the amount of chlorophyll *a* and pheophytin *a* in marine and freshwater algae by visible spectrophotometry. Uncorrected chlorophyll *a* is calculated using the trichromatic equation. Corrected chlorophyll *a* and pheophytin are calculated using the monochromatic equation. The absorption-peak-ratio (chlorophyll/pheophytin) is also determined.
- 3) Preservation Method: Whole water is collected in brown Nalgene bottles, placed on ice in the dark, and delivered to the FLDEP lab within 36 hours for filtration.

g) Parameter: TSS

- 1) Method Reference: This method is based on Standard Methods 2540 D-1997.
- 2) Method Description: A well-mixed sample is filtered through a pre-weighed glass fiber filter. The filter and any residue are then dried to a constant weight at 103-105 °C. The filter is cooled in a desiccator, weighed and the result used to compute the TSS of the sample.
- 3) Preservation Method: Whole water is placed on ice in the dark for analysis within 7 days.

14) Field and Laboratory QA/QC programs:

a) Precision

- i) Field Variability – Field Blanks are included in all monthly sampling events. ANERR staff collected field triplicate samples from a successive grab sample. Triplicate samples are collected from separate grabs at one sampling station every other month, rotating through stations. There were no field triplicates collected during diel sampling.
- ii) Laboratory Variability – Method blanks (MB) and duplicate samples are run with every sample batch. Batches are groups of 20 or less samples that are analyzed concurrently. Precision is measured by Relative Percent Difference (RPD).
- iii) Inter-organizational splits – None.

b) Accuracy

- i) Sample Spikes – At least two sample spikes are performed with each sample batch. The acceptance limits for sample or spike duplicates is a RPD of less than 20% if both results are above the PQL. Laboratory fortified blanks are run with each sample batch, acceptance limits for recovery are 85-115%.
- ii) Standard Reference Material Analysis – Check standards are included in each batch and at the beginning and end of each run. Check standard acceptance limits are 85-115% recovery. (FLDEP Central Laboratory NU-043-2.16)
- iii) Cross Calibration Exercises – None for 2013.

15) QAQC flag definitions:

QAQC flags provide documentation of the data and are applied to individual data points by insertion into the parameter's associated flag column (header preceded by an F_). QAQC flags are applied to the nutrient data during secondary QAQC to indicate data that are out of sensor range low (-4), rejected due to QAQC checks (-3), missing (-2), optional and were not collected (-1), suspect (1), and that have been corrected (5). All remaining data are flagged as having passed initial QAQC checks (0) when the data are uploaded and assimilated into the CDMO ODIS as provisional plus data. The historical data flag (4) is used to indicate data that were submitted to the CDMO prior to the initiation of secondary QAQC flags and codes (and the use of the automated primary QAQC system for WQ and MET data). This flag is only present in historical data that are exported from the CDMO ODIS.

- 4 Outside Low Sensor Range
- 3 Data Rejected due to QAQC
- 2 Missing Data
- 1 Optional SWMP Supported Parameter
- 0 Data Passed Initial QAQC Checks
- 1 Suspect Data
- 4 Historical Data: Pre-Auto QAQC
- 5 Corrected Data

16) QAQC code definitions:

QAQC codes are used in conjunction with QAQC flags to provide further documentation of the data and are also applied by insertion into the associated flag column. There are three (3) different code categories, general, sensor, and comment. General errors document general problems with the sample or sample collection, sensor errors document common sensor or parameter specific problems, and comment codes are used to further document conditions or a problem with the data. Only one general or sensor error and one comment code can be applied to a particular data point. However, a record flag column (F_Record) in the nutrient data allows multiple comment codes to be applied to the entire data record.

General errors

- GCM Calculated value could not be determined due to missing data
- GCR Calculated value could not be determined due to rejected data
- GDM Data missing or sample never collected
- GQD Data rejected due to QA/QC checks
- GQS Data suspect due to QA/QC checks
- GSM See metadata

Sensor errors

- SBL Value below minimum limit of method detection
- SCB Calculated value could not be determined due to a below MDL component
- SCC Calculation with this component resulted in a negative value
- SNV Calculated value is negative
- SRD Replicate values differ substantially
- SUL Value above upper limit of method detection

Parameter Comments

CAB	Algal bloom
CDR	Sample diluted and rerun
CHB	Sample held beyond specified holding time
CIP	Ice present in sample vicinity
CIF	Flotsam present in sample vicinity
CLE	Sample collected later/earlier than scheduled
CRE	Significant rain event
CSM	See metadata
CUS	Lab analysis from unpreserved sample

Record comments

CAB	Algal bloom
CHB	Sample held beyond specified holding time
CIP	Ice present in sample vicinity
CIF	Flotsam present in sample vicinity
CLE	Sample collected later/earlier than scheduled
CRE	Significant rain event
CSM	See metadata
CUS	Lab analysis from unpreserved sample

Cloud cover

CCL	clear (0-10%)
CSP	scattered to partly cloudy (10-50%)
CPB	partly to broken (50-90%)
COC	overcast (>90%)
CFY	foggy
CHY	hazy
CCC	cloud (no percentage)

Precipitation

PNP	none
PDR	drizzle
PLR	light rain
PHR	heavy rain
PSQ	squally
PFQ	frozen precipitation (sleet/snow/freezing rain)
PSR	mixed rain and snow

Tide stage

TSE	ebb tide
TSF	flood tide
TSH	high tide
TSL	low tide

Wave height

WH0	0 to <0.1 meters
WH1	0.1 to 0.3 meters
WH2	0.3 to 0.6 meters
WH3	0.6 to > 1.0 meters

WH4	1.0 to 1.3 meters
WH5	1.3 or greater meters
<i>Wind direction</i>	
N	from the north
NNE	from the north northeast
NE	from the northeast
ENE	from the east northeast
E	from the east
ESE	from the east southeast
SE	from the southeast
SSE	from the south southeast
S	from the south
SSW	from the south southwest
SW	from the southwest
WSW	from the west southwest
W	from the west
WNW	from the west northwest
NW	from the northwest
NNW	from the north northwest
<i>Wind speed</i>	
WS0	0 to 1 knot
WS1	> 1 to 10 knots
WS2	> 10 to 20 knots
WS3	> 20 to 30 knots
WS4	> 30 to 40 knots
WS5	> 40 knots

17) Other remarks/notes:

Data may be missing due to problems with sample collection or processing. Laboratories in the NERRS System submit data that are censored at a lower detection rate limit, called the Method Detection Limit or MDL. MDLs for specific parameters are listed in the Laboratory Methods and Detection Limits Section (Section II, Part 12) of this document. Concentrations that are less than this limit are censored with the use of a QAQC flag and code, and the reported value is the method detection limit itself rather than a measured value. For example, if the measured concentration of NO₃F was 0.0005 mg/l as N (MDL=0.0008), the reported value would be 0.0008 and would be flagged as out of sensor range low (-4) and coded SBL. In addition, if any of the components used to calculate a variable are below the MDL, the calculated variable is removed and flagged/coded -4 SCB. If a calculated value is negative, it is rejected and all measured components are marked suspect. If additional information on MDL's or missing, suspect, or rejected data is needed, contact the Research Coordinator at the Reserve submitting the data.

Note: The way below MDL values are handled in the NERRS SWMP dataset was changed in November of 2011. Previously, below MDL data from 2007-2010 were also flagged/coded, but either reported as the measured value or a blank cell. Any 2007-2011 nutrient/pigment data downloaded from the CDMO prior to November of 2011 will reflect this difference.

01/10/2014 11:45 apaesnut: Low winter tides delayed deployment, which in turn delayed retrieval of the ISCO. Retrieval after the last scheduled sample at 11:45 would have made it too late to process and transport samples to the DEP lab, therefore the ISCO was retrieved prior to the last sample being collected.

03/04/2014 09:45 – 03/05/2014 10:45 apaesnut; Diel samples were not iced during transport to FLDEP laboratory. All data are rejected.

03/04/2014 12:16 apadbnut; CHLA_N, uncCHLA_N, and PHEA sample dropped in laboratory, no analysis performed.

06/10/2014 09:30 – 06/11/2014 08:00 apaesnut; Automated ISCO sampler collected short sample volumes. No samples were collected for CHLA_N, uncCHLA_N, PHEA, or TSS.

6/11/2014 11:30 apaesnut; Sample is missing, ISCO was retrieved early due to weather.

12/02/2014 10:33 apaegnut; NH4_F sample discarded by laboratory prior to analysis.

12/02/2014 10:15 apaesnut; NH4_F and NO23_F samples were not collected.

12/02/2014 and 12/03/2014: Weather issues forced sampling to be conducted over two days. No samples were collected at these sites: apanhnut, apapcnut, apawpnut.

Sample holding times are available upon request.