

Tijuana River (TR) NERR Nutrient Metadata
January to December 2024
Latest Update: July 1, 2025

Note: This is a provisional metadata document; it has not been authenticated as of its download date. Contents of this document are subject to change throughout the QAQC process and it should not be considered a final record of data documentation until that process is complete. Contact the CDMO (cdmosupport@baruch.sc.edu) or reserve with any additional questions.

I. Data Set and Research Descriptors

1) Principal investigator(s) and contact persons –

Jeff Crooks, Research Coordinator

301 Caspian Way
Imperial Beach, CA 91932
Phone: (619) 575-3613 x333
Fax: (619) 575-6913
jcrooks@tijuanaestuary.org

Monica Almeida, Research Associate

301 Caspian Way
Imperial Beach, CA 91932
Phone: (619) 575-3613 x322
malmeida@trnerr.org

Justin McCullough, Research Associate

301 Caspian Way
Imperial Beach, CA 91932
Phone: (619) 575-3613 x323
jmccullough@trnerr.org

Stephany Garcia, Research Associate

301 Caspian Way
Imperial Beach, CA 91932
Phone: (619) 575-3613 x321
sgarcia@trnerr.org

Susan M. Becker, Laboratory Manager

Scripps Institution of Oceanography,
University of California, San Diego
9500 Gilman Dr. #0236
La Jolla, CA 92093
sio-sts-chemists@ucsd.edu

2) Research objectives –

The Tijuana River National Estuarine Research Reserve (TRNERR) represents the largest, most intact coastal marsh system remaining in Southern California. It has contiguous beach, dune, tidal channel, mudflat, marsh, transitional, and upland habitat. It is also home to numerous threatened and endangered species. Because of its highly urbanized setting, situated between the cities of Tijuana, Baja California, Mexico, and San Diego, California, USA, it is heavily impacted. A primary management concern is transboundary flows of the Tijuana River, which convey anthropogenic pollutants (primarily associated with partially-treated and untreated wastewater), nutrients, and sediment. About a quarter of the reserve's 2,531 acres are tidally influenced and few channels are deep enough for datalogger deployment. Currently, there are two SWMP stations located within the TRNERR boundaries, and two SWMP stations are located nearby in south San Diego Bay. Station locations are designed to investigate spatial patterns of water quality parameters, with comparisons between the Tijuana Estuary and San Diego Bay. In addition, telemetry of Tijuana Estuary water quality stations informs management action, particularly related to potential closure of the tidal inlet by wave-driven accumulation of sediment. Mouth closures (detected by cessation of tidal action as indicated with the water level sensors on the dataloggers) can cause anoxia, mortality of fish and shellfish, and flooding.

Two stations were originally set up: a “control” station Oneonta Slough (OS), which is still in place, was established on the northern end of Oneonta Slough, relatively far away from the main source of river-borne pollution. Another station, River Channel (RC), was situated in a site most affected by sewage outflow. Datalogger deployment at RC, however, was continually interrupted by both shifting sediment and massive wracks of kelp (*Macrocystis pyrifera*), which would often bury the deployment set-up on incoming tides. After a number of different deployment designs were implemented without success, data collection at the RC site was terminated in 2004. Another station was located at the inlet to the Model Marsh (MM), a constructed 20-acre restoration site in the southern arm of the estuary. The Model Marsh was opened to tidal flushing in February 2000 and datalogging at the station began in October 2000. The site was discontinued in January 2008, again due to heavy sedimentation. The Boca Rio (BR) site is located near the mouth of the Tijuana River, although the mouth has migrated south in recent years. The BR station was established in December 2004 to replace the RC station, and remains active and is the site closest to the interface of the river and ocean.

The South Bay (SB) station was established in January 2008 and is located at the mouth of Otay River, which flows into South San Diego Bay. It is within the San Diego National Wildlife Refuge Complex, which also includes portions of the TRNERR. It was established to document conditions in the south bay, particularly associated with a marsh restoration in the adjacent salt ponds. The Pond Eleven (PE) station was located in a non-tidal salt pond adjacent to the South Bay logger. A tide gate was the only source of water into the pond, which was one of the first in a series of ponds with increasing salinities. The PE sonde was deployed from July 2008 to September 2010. The US Fish and Wildlife Service began restoration of this area, including Pond Eleven, from September 2010 to its completion in October 2011. A levee was breached to open Pond Eleven to the bay, which made the area tidal, and channels were excavated to further enhance circulation. Due to extensive restoration, the site had to be relocated. Datasonde deployments and nutrient data collection began in January 2012 at a new location site named Pond Restored (PR) (the name change was warranted because of the different location and profound differences in the pond before and after restoration). The Pond Restored datalogger is located approximately 560 meters southwest from where the Pond Eleven datalogger was originally. The images below show pre- and post-restoration of the salt ponds and the station locations. The post restoration photo includes the PE site as a reference to the new PR site. No sampling currently occurs at the PE site.



a) Monthly grab sampling program

Grab samples were collected monthly at all SWMP water quality sonde sites. Monthly grab sampling provides information as to the spatial differences in nutrient levels within the system. The placement of these sites also offers the opportunity to better isolate the possible origin of nutrient inputs as well as the degree of tidal flushing.

b) Diel sampling program (mention if samples were taken over a lunar day)

Diel samples were collected monthly at one site. This sampling method provides a temporal perspective of nutrient flux over one tidal cycle at this particular site. Given the location of the site, the diel sampling also provides valuable information as to the degree of tidal flushing of nutrients from the estuary.

3) Research methods –

All samples were collected in compliance with the NERRS Nutrient and Chlorophyll Monitoring SOP version 1.7.

Prior to sampling, all sample containers (ISCO 1L Polyethylene bottles for diel samples and Nalgene 1L wide mouth amber opaque HDPE bottles for grab samples) were acid washed. The washing procedure consisted of four rinses with DI water, a gentle squirt of Liquinox phosphate free detergent, six rinses with DI water, a soak of ten to twenty minutes in 10% HCl, and, lastly, six rinses with DI water.

a) Monthly grab sampling program

Monthly grab samples were taken at the SWMP sites (Boca Rio, Oneonta Slough, Pond Restored, and South Bay). All samples were collected less than three hours prior to the projected slack low tide ending a diel sampling cycle. Samples were collected by hand near the surface in a vertical motion, using amber wide mouth 1 L acid washed (10% HCl) Nalgene bottles. Sample bottles were rinsed three times with ambient water before sampling. Sampling typically followed a dry period of 72 hours or greater, unless otherwise noted (see Other Remarks at the end of this document). Duplicate samples were taken at each site sequentially, within one minute of each other and adjacent to the datasonde deployed at each station.

At the time of nutrient sampling, in-situ measurements of temperature, specific conductivity, salinity and dissolved oxygen were taken using an YSI Digital Professional Series water quality handheld instrument. When handheld instrument was not available, nutrient samples were taken as close as possible to a 15-minute mark, when a water quality measurement is taken by the adjacent datasonde. These measurements are included in the datasheet.

b) Diel sampling program

Diel samples were collected at the Boca Rio site. Samples were taken at intervals of 2hrs 15mins for a total of, at least, 11 samples over one lunar day. Samples were collected using an ISCO model 6712 autosampler with a sixty-foot sample tube and polypropylene strainer. The strainer was suspended in a minnow trap (approximately 25 cm above the channel bottom) within 20m of the associated datasonde. The sampler was located approximately 1.5m above the strainer on the channel's bank. The sampler was programmed to collect 0.5 liter per sample utilizing ISCO's 24-bottle kit in a standard tub stocked with ice. At the time of retrieval, the sampling program is suspended, the sample bottles are capped and, typically, reach the reserve's lab within one hour for filtration.

After filtration, all samples were delivered, generally, within 72 hours to the Chemistry Lab at the Oceanographic Data Facility, Scripps Institution of Oceanography (SIO), University of California, San Diego. Samples were frozen at -20°C and analyzed within 28 days (30 days for CHLA) unless indicated in red font in the "Samples collection and analysis date" table (see Other Remarks). The Oceanographic Data Facility's nutrient analysis procedures can be found at <https://scripps.ucsd.edu/ships/shipboard-technical-support/odf/documentation/nutrient-analysis>.

In addition to required parameters, the TRNERR also began reporting silicate in 2015. Chlorophyll analyses were performed in house at the TRNERR until June 2017. Starting in August 2017, the frozen chlorophyll samples were also delivered to the SIO lab. The chlorophyll analysis method used by SIO lab can be found at <http://calcofi.org/ccpublications/calcofi-methods/8-chlorophyll-methods.html>.

4) Site location and character –

The four SWMP sites sit in the southwest corner of San Diego County, just north of the US / Mexico border. Two are located in the Tijuana River Estuary, at the terminus of the multinational (US, Mexico, and indigenous Kumeyaay) Tijuana River watershed. Two are associated with the adjacent watershed of the Otay River in the southern end of San Diego Bay, which has been highly modified by channelization of the river and creation of a salt production facility in the bay, which dates back to the 1870's. The Tijuana River Estuary and San Diego Bay are both in the Silver Strand Littoral Cell, with their barrier beaches formed by sediment input from the Tijuana River. Historically, the Tijuana River Estuary and San Diego Bay were part of a large coastal wetland complex, separated by non-tidal wetland in what is now the City of Imperial Beach.

All estuaries in the region reside in a Mediterranean-climate, and are characterized as low-inflow estuaries, with natural freshwater input largely confined to the rainy season (October – March). Urbanization of watersheds have led to perennialization of many formerly ephemeral streams. In the Tijuana River, flows are typically contaminated with sewage, and infrastructure in the US and Mexico has been built to try and manage these flows. The South Bay International Wastewater Plant began operation in 1999, and was designed to capture dry-weather flows. Any rain or infrastructure failures in Mexico typically exceed the capacity of the plant and result in transboundary flows of contaminated water. Persistent transboundary flows in to the Tijuana River Estuary have been especially problematic since 2022.

Tidal exchange at the inlet of the Tijuana Estuary is limited by an intertidal sill, with occasional mouth closures. Excessive sedimentation in the Tijuana River Estuary has greatly decreased tidal prism, which has exacerbated mouth closures. The South San Diego Bay sites are fully tidal. Both the Tijuana River Estuary and south San Diego Bay are characterized by coastal salt marsh, including cordgrass (*Spartina foliosa*) and pickleweed (*Salicornia pacifica*). Both are also sites of ongoing restoration programs aimed at recovering lost salt marsh habitat.

Site name	Boca Rio (BR)
Latitude and longitude	32° 33' 33.7" N, 117° 7' 44.3" W
Tidal range (<i>meters</i>)	Approx. 1.6, but variable depending on the height of the intertidal sill at the inlet, which limits the extent of low tides in the estuary compared to the open coast.
Salinity range (<i>psu</i>)	1 (extreme rain events) to 36 (average of 33)
Type and amount of freshwater input	The Tijuana River Estuary is low-inflow, and the dominant freshwater source is the Tijuana River. Stream flows in the river vary considerably from season to season and year to year. The naturally-ephemeral Tijuana River now has year-round flow upstream in Tijuana, Mexico, although wastewater infrastructure (when operational) diverts low flows (less than 1 cms) and prevents anthropogenic surface flows from reaching the estuary. Additional freshwater sources are storm drains located mostly in the northern arm of the estuary from runoff from the adjacent military airfield and residential area.
Water depth (<i>meters, MLW</i>)	1.3 (estimated)
Sonde distance from bottom (<i>meters</i>)	Approx. 0.25
Bottom habitat or type	Silt and clay, some sand
Pollutants in area	Freshwater discharge with sewage

Description of watershed	The Tijuana River watershed is one of the largest in the region, at 4500 km ² , approx. 75% of which is in Mexico. The lower portion is heavily urbanized, with the channelized river flowing through the city of Tijuana, Mexico, before entering the United States.
--------------------------	--

Site name	Oneonta Slough (OS)
Latitude and longitude	32° 34' 6.0" N, 117° 7' 52.6" W
Tidal range (<i>meters</i>)	Approx. 1.6, but variable depending on the height of the intertidal sill at the inlet, which limits the extent of low tides in the estuary compared to the open coast.
Salinity range (<i>psu</i>)	1 (extreme rain events) to 39 (average of 32 ppt)
Type and amount of freshwater input	The Tijuana River Estuary is low-inflow, and the dominant freshwater source is the Tijuana River. Stream flows in the river vary considerably from season to season and year to year. The naturally-ephemeral Tijuana River now has year-round flow upstream in Tijuana, Mexico, although wastewater infrastructure (when operational) diverts low flows (less than 1 cms) and prevents anthropogenic surface flows from reaching the estuary. Additional freshwater sources are storm drains located mostly in the northern arm of the estuary from runoff from the adjacent military airfield and residential area.
Water depth (<i>meters, MLW</i>)	1.1 (estimated)
Sonde distance from bottom (<i>meters</i>)	Approx. 0.25
Bottom habitat or type	Silt and clay
Pollutants in area	Freshwater discharge with sewage, runoff from streets.
Description of watershed	The Tijuana River watershed is one of the largest in the region, at 4500 km ² , approx. 75% of which is in Mexico. The lower portion is heavily urbanized, with the channelized river flowing through the city of Tijuana, Mexico, before entering the United States.

Site name	Pond Restored (PR)
Latitude and longitude	32° 35' 45.9", 117° 7' 5.5" W
Tidal range (<i>meters</i>)	Approx. 2.7
Salinity range (<i>psu</i>)	4 (extreme rain event) to 39 (average of 35)
Type and amount of freshwater input	Highly seasonal flows from the Otay River, which enters into the extreme south end of San Diego Bay. Salinity can also be affected by occasional leakage from high-salinity ponds of the South Bay Salt Works.
Water depth (<i>meters, MLW</i>)	1.6 (estimated)

Sonde distance from bottom (<i>meters</i>)	Approx. 0.25
Bottom habitat or type	Silt and clay
Pollutants in area	Legacy metal and synthetic organics in San Diego Bay.
Description of watershed	The 40 km-long Otay River originates in the mountains of southern San Diego County, and is dammed at the Otay Reservoir. The watershed is 410km ² , and the lower watershed includes the City of Chula Vista, California.

Site name	South Bay (SB)
Latitude and longitude	32° 36' 3.6" N, 117° 06' 57.0" W
Tidal range (<i>meters</i>)	Approx. 2.7
Salinity range (<i>psu</i>)	4 (extreme rain event) to 39 (average of 35)
Type and amount of freshwater input	Highly seasonal flows from the Otay River, which enters into the extreme south end of San Diego Bay. Salinity can also be affected by occasional leakage from high-salinity ponds of the South Bay Salt Works.
Water depth (<i>meters, MLW</i>)	1.8 (estimated)
Sonde distance from bottom (<i>meters</i>)	Approx. 0.25
Bottom habitat or type	Silt and clay
Pollutants in area	Legacy metal and synthetic organics in San Diego Bay.
Description of watershed	The 40 km-long Otay River originates in the mountains of southern San Diego County, and is dammed at the Otay Reservoir. The watershed is 410km ² , and the lower watershed includes the City of Chula Vista, California.

All Tijuana River NERR historical nutrient/pigment monitoring stations:

Station Code	SWMP Status	Station Name	Location	Active Dates	Reason Decommissioned	Notes
tjrbrnut	P	Boca Rio	32° 33' 33.70 N, 117° 7' 44.30 W	12/01/2004 - current	NA	Grab sample until present; Diel sample from 1/18/2007 to 12/6/2012 and from 1/20/2016 to present
tjrosnut	P	Oneonta Slough	32° 34' 6.00 N, 117° 7' 52.60 W	09/01/2002 - current	NA	Grab sample until present; diel sample

						from 9/1/2002 until 12/06/2006
tjrprnut	P	Pond Eleven Restored	32° 35' 45.90 N, 117° 07' 5.50 W	01/01/2012 - current	NA	Grab sample until present; diel sample from 1/8/2013 to 12/18/2015
tjrbsnut	P	South Bay	32° 36' 3.60 N, 117° 6' 57.00 W	01/01/2008 - current	NA	Grab sample
tjrpenut	P	Pond Eleven	32° 36' 3.54 N, 117° 6' 58.46 W	07/01/2008 – 12/31/2011	Sampling at this site was interrupted due to an extensive Restoration project	Grab sample; restoration project was concluded in October 2011. Pond Eleven site was relocated and renamed to Pond Eleven Restored. Sampling resumed in January 2013.
tjrmmnut	P	Model Marsh	32° 32' 52.08 N, 117° 7' 22.80 W	09/01/2002 – 07/30/2008	Heavy sedimentation compromised the datalogger station	Grab sample
tjrtnut	P	Tidal Linkage	32° 34' 27.84 N, 117° 7' 37.92 W	09/01/2002 – 12/06/2007	Site too shallow for diel samples; heavy sedimentation compromised the datalogger station	Grab sample; diel sample on 7/27/2006
tjrrenut	P	River Channel	32° 33' 28.08 N, 117° 6' 21.96 W	09/01/2002 – 11/30/2004	Heavy sedimentation compromised the datalogger station	Grab sample; replaced by Boca Rio site
tjrhpnut	NA	Helicopter Pad	32° 33' 31.46 N, 117° 7' 9.34 W	09/01/2002 - 09/30/2002	Grab sample collected to test site suitability	Data only available by contacting the reserve directly
tjrmmnut	NA	River Mouth	32° 33' 13.32 N, 117° 7' 34.32 W	09/01/2002 – 12/01/2002	Grab samples collected to test site feasibility	Data only available by contacting the reserve directly

5) Coded variable definitions –

Station code names:

tjrbrnut = Tijuana River Reserve Boca Rio nutrients
tjrosnut = Tijuana River Reserve Oneonta Slough nutrients
tjrbsnut = Tijuana River Reserve South Bay nutrients
tjrpenut = Tijuana River Reserve Pond Restored nutrients

monthly grab sample program = 1
diel grab sample program = 2

**6) Data collection period – Grab
Samples**

Boca Rio (BR)

Station code	Date	Rep1	Rep 2
tjrbrnut	1/24/2024	13:50	13:51
tjrbrnut	2/22/2024	14:27	14:28
tjrbrnut	3/19/2024	12:33	12:34
tjrbrnut	4/17/2024	10:47	10:48
tjrbrnut	5/16/2024	11:15	11:16
tjrbrnut	6/19/2024	11:01	11:02
tjrbrnut	7/23/2024	14:12	14:13
tjrbrnut	8/19/2024	13:57	13:58
tjrbrnut	9/16/2024	13:27	13:28
tjrbrnut	10/16/2024	12:00	12:01
tjrbrnut	11/13/2024	13:30	13:31
tjrbrnut	12/11/2024	11:53	11:54

Oneonta Slough (OS)

Station code	Date	Rep1	Rep 2
tjrosnut	1/24/2024	14:33	14:34
tjrosnut	2/22/2024	15:00	15:01
tjrosnut	3/19/2024	14:20	14:21
tjrosnut	4/17/2024	10:17	10:18
tjrosnut	5/16/2025	11:50	11:51
tjrosnut	6/19/2024	11:34	11:35
tjrosnut	7/23/2024	13:40	13:41
tjrosnut	8/19/2024	13:00	13:01
tjrosnut	9/17/2024	11:57	11:58
tjrosnut	10/16/2024	13:00	13:01
tjrosnut	11/13/2024	14:00	14:01
tjrosnut	12/11/2024	11:28	11:29

Pond Restored (PR)

Station code	Date	Rep1	Rep 2
tjrprnut	1/24/2024	13:23	13:24
tjrprnut	2/21/2024	12:26	12:27
tjrprnut	3/19/2024	11:44	11:45
tjrprnut	4/17/2024	11:30	11:31
tjrprnut	5/16/2024	8:18	8:19
tjrprnut	6/19/2024	11:50	11:51
tjrprnut	7/23/2024	14:30	14:31
tjrprnut	8/20/2024	13:41	13:42
tjrprnut	9/17/2024	12:36	12:37
tjrprnut	10/16/2024	12:15	12:16

tjrprnut	11/13/2024	11:46	11:47
tjrprnut	12/11/2024	11:27	11:28

South Bay (SB)

Station code	Date	Rep1	Rep 2
tjrbsnut	1/24/2024	12:34	12:35
tjrbsnut	2/21/2024	11:30	11:31
tjrbsnut	3/19/2024	12:09	12:10
tjrbsnut	4/17/2024	10:19	10:20
tjrbsnut	5/16/2024	8:47	8:48
tjrbsnut	6/19/2024	11:05	11:06
tjrbsnut	7/23/2024	14:00	14:01
tjrbsnut	8/20/2024	13:09	13:10
tjrbsnut	9/17/2024	13:10	13:11
tjrbsnut	10/16/2024	12:45	12:46
tjrbsnut	11/13/2024	12:23	12:24
tjrbsnut	12/11/2024	10:59	11:00

Diel Samples

Boca Rio (BR)

Station code	Start Date	Start Time	End Date	End Time
tjrbrnut	1/23/2024	12:15	1/24/2024	10:45
tjrbrnut	2/20/2024	11:30	2/21/2024	10:00
tjrbrnut	3/18/2024	13:00	3/19/2023	11:30
tjrbrnut	4/16/2024	12:00	4/17/2024	10:30
tjrbrnut	5/15/2024	11:00	5/16/2024	9:30
tjrbrnut	6/18/2024	9:45	6/19/2024	8:15
tjrbrnut	7/22/2024	15:15	7/23/2024	13:45
tjrbrnut	8/19/2024	12:15	8/20/2024	10:45
tjrbrnut	9/16/2024	13:30	9/17/2024	12:00
tjrbrnut	10/15/2024	13:15	10/16/2024	11:45
tjrbrnut	11/12/2024	12:30	11/13/2024	11:00
tjrbrnut	12/10/2024	11:30	12/11/2024	10:00

Note that the time stamps are based on a 24-hour clock and refer to Pacific Standard Time. The date format is mm/dd/yyyy. Water quality parameters such as Specific Conductivity, Salinity, Dissolved Oxygen Saturation and Temperature are measured by YSI Professional Plus water quality instrument upon grab samples collection. These measurements are included in the nutrient datasheet.

7) Associated researchers and projects—

The research program at the TRNERR focuses on adaptive approaches to wetlands management, which involves coupling scientific investigation with management action. One focal area of research continues to be adaptive restoration, and the TRNERR has a long history of science-based restoration efforts. These programs incorporate descriptive and experimental approaches to investigate biotic and abiotic responses to marsh restoration, including

ways to better achieve desired ecosystem responses. Two SWMP sites, based in South San Diego Bay, are associated with a marsh restoration in that area. Another active area of research is invasive species ecology and management. Although estuaries are typically invaded by a broad suite of species from many habitat types, current research is focusing on terrestrial and riparian invaders able to cross ecotones and invade salt marsh habitats. Researchers at the TRNERR are investigating mechanisms of invasions, impacts of invaders, and ecosystem recovery after exotic species control.

NERR SWMP water quality and weather data are used in a variety of reserve-based and external research and education programs. Water quality data from the Tijuana River, which rarely experiences mouth closures, provides an interesting contrast to data from other regional systems that experience frequent closure events. Also, SWMP water quality data are incorporated into a high school curriculum developed at the reserve. Tier 1 nutrient sampling is being conducted at all water quality datalogger stations. NERR SWMP meteorological sampling is being conducted at 1 station which is located near the former Tidal Linkage water quality station. In addition, much of the reserve is used as a test bed for research related to adaptive marsh restoration. NERR SWMP WQ and MET data are collected every 15-minutes and may be correlated with this nutrient/pigment dataset. These data are available at www.nerrsdata.org.

8) Distribution –

NOAA retains the right to analyze, synthesize and publish summaries of the NERRS System-wide Monitoring Program data. The NERRS retains the right to be fully credited for having collected and processed the data. Following academic courtesy standards, the NERR site where the data were collected should be contacted and fully acknowledged in any subsequent publications in which any part of the data are used. The data set enclosed within this package/transmission is only as good as the quality assurance and quality control procedures outlined by the enclosed metadata reporting statement. The user bears all responsibility for its subsequent use/misuse in any further analyses or comparisons. The Federal government does not assume liability to the Recipient or third persons, nor will the Federal government reimburse or indemnify the Recipient for its liability due to any losses resulting in any way from the use of this data.

Requested citation format:

NOAA National Estuarine Research Reserve System (NERRS). System-wide Monitoring Program. Data accessed from the NOAA NERRS Centralized Data Management Office website: www.nerrsdata.org; accessed 12 October 2024.

NERR nutrient data and metadata can be obtained from the Research Coordinator at the individual NERR site (please see Principal investigators and contact persons), from the Data Manager at the Centralized Data Management Office (please see personnel directory under the general information link on the CDMO home page) and online at the CDMO home page www.nerrsdata.org. Data are available in comma separated version format.

II. Physical Structure Descriptors

9) Entry verification –

TJR NERR staff collected, filtered and delivered the samples to the chemistry lab at the Oceanographic Data Facility, Scripps Institution of Oceanography, University of California, San Diego, where lab analyses were performed. The results are reported in $\mu\text{M/L}$. For purposes of consistency in the NERR System, the TJR NERR calculates the concentrations as mg/l based on atomic weights of 14.01, 30.97, and 28.09 for N, P, and Si, respectively. Therefore, TJR NERR multiplies the concentrations reported by 0.01401, 0.03097, and 0.02809 to yield concentrations in mg/L as N, P, and Si, respectively.

Nutrient data are entered into a Microsoft Excel worksheet and processed using the NutrientQAQC Excel macro. The NutrientQAQC macro sets up the data worksheet, metadata worksheets, and MDL worksheet; adds chosen parameters and facilitates data entry; allows the user to set the number of significant figures to be reported for each parameter and rounds using banker's rounding rules; allows the user to input MDL values and then automatically flags/codes measured values below MDL and inserts the MDL; calculates parameters chosen by the user and

automatically flags/codes for component values below MDL, negative calculated values, and missing data; allows the user to apply QAQC flags and codes to the data; produces summary statistics; graphs selected parameters for review; and exports the resulting data file to the CDMO for tertiary QAQC and assimilation into the CDMO's authoritative online database.

Data entry and verification was performed by Monica Almeida (please see **Principal investigators and contact persons**).

10) Parameter titles and variable names by category –

Required NOAA NERRS System-wide Monitoring Program nutrient parameters are denoted by an asterisk “*”.

Data Category	Parameter	Variable Name	Units of Measure
Phosphorus and Nitrogen:			
	*Orthophosphate	PO4F	mg/L as P
	*Ammonium, Filtered	NH4F	mg/L as N
	*Nitrite + Nitrate, Filtered	NO23F	mg/L as N
	Dissolved Inorganic Nitrogen	DIN	mg/L as N
Plant Pigments:			
	*Chlorophyll a	CHLA_N	µg/L
Carbon:			
Other Lab Parameters:			
	Silicate, Filtered	SiO4F	mg/L as SI
Field Parameters: (Grab samples):			
	Water Temperature	WTEM_N	°C
	Specific Conductivity	SCON_N	mS/cm
	Salinity	SALT_N	ppt
	Dissolved Oxygen	DO_N	mg/L
	Dissolved Oxygen Saturation	DO_S_N	%

Notes:

1. Time is coded based on a 2400 clock and is referenced to Standard Time.
2. Reserves have the option of measuring either NO₂ and NO₃ or they may substitute NO₂₃ for individual analyses if they can show that NO₂ is a minor component relative to NO₃. NO₂ levels are predominately low at TJR Reserve, and individual analysis results have been measured but not reported. Results of individual analyses for NO₂ are available upon request from the Research Coordinator at the specific NERR site. Please refer to the Principal investigators and contact persons for more information.

11) Measured or calculated laboratory parameters –

a) Parameters measured directly

Nitrogen species:	NH4F, NO23F
Phosphorus species:	PO4F
Other:	CHLA_N, SiO4F

b) Calculated parameters

DIN	NO23F+NH4F
-----	------------

12) Limits of detection –

Method detection limit as defined in the Code of Federal Regulations, 40 CFR 136, Appendix B, as: “the minimum concentration of a substance that can be measured and reported with 99% confidence that the analyte concentration is greater than zero and is determined from analysis of a sample in a given matrix containing the analyte.”

MDLs were reported by the Chemistry Lab at the Oceanographic Data Facility, Scripps Institution of Oceanography, University of California, San Diego. The lab determines MDLs by running a minimum of seven samples prepared at low concentrations, and multiplying the calculated standard deviation by 3 (approximately, this factor depends on the number of measurements). The lab MDL SOP is based on the methodology described in the “**Definition and Procedure for the Determination of the Method Detection Limit, Revision 2**” by the U.S. Environmental Protection Agency, which can be found at:

https://www.epa.gov/sites/production/files/2016-12/documents/mdl-procedure_rev2_12-13-2016.pdf

These values are revisited annually.

<u>Parameter</u>	<u>Start Date</u>	<u>End Date</u>	<u>MDL</u>	<u>Revisited</u>
PO4F	1/1/2024	6/25/2024	0.0004	7/28/2023
PO4F	6/26/2024	12/31/2024	0.0002	6/26/2024
NH4F	1/1/2024	6/25/2024	0.002	7/28/2023
NH4F	6/26/2024	12/31/2024	0.0017	6/26/2024
NO23F	1/1/2024	6/25/2024	0.0009	7/28/2023
NO23F	6/26/2024	12/31/2024	0.0008	6/26/2024
CHLA_N	1/1/2024	1/31/2024	0.0003	1/24/2024
CHLA_N	2/1/2024	6/25/2024	0.0009	8/9/2023
CHLA_N	6/26/2024	12/31/2024	0.0005	6/26/2024
SiO4F	1/1/2024	6/25/2024	0.005	7/28/2023
SiO4F	6/26/2024	12/31/2024	0.021	6/26/2024

13) Laboratory methods –

[

a) **Parameter: PO4F**

Method Reference: Orthophosphate is analyzed using a modification of the Bernhardt and Wilhelms method. Bernhardt, H., and Wilhelms, A., 1967. "The continuous determination of low level iron, soluble phosphate and total phosphate with the AutoAnalyzer," Technicon Symposia, I, pp.385-389.

Method Descriptor: Acidified ammonium molybdate is added to a water sample to produce phosphomolybdic acid, which is then reduced to phosphomolybdous acid (a blue compound) following the addition of dihydrazine sulfate. The sample is passed through a 10mm flowcell and absorbance is measured at 820nm. The absorbance is proportional to the concentration of orthophosphate in the sample.

Preservation Method: Filtered through both Whatman glass fiber filter (.7µm pore size) and MF-Millipore (.45µm pore size) membrane filter, and stored at or below -20°C.

Instrumentation: AA3 Autoanalyzer made my SEAL Analytical.

b) **Parameter: NH4F**

Method Reference: Ammonia is analyzed using the method described by Kerouel and Aminot. Kerouel, R. and Aminot, A., 1997. "Fluorometric determination of ammonia in sea and estuarine waters by direct segmented flow analysis." Marine Chemistry, vol 57, no. 3-4, pp. 265-275.

Method Descriptor: The sample is combined with a working reagent made up of ortho-phthalaldehyde, sodium sulfite and borate buffer and heated to 75°C. Fluorescence proportional to the ammonia concentration is emitted at 460nm following excitation at 370nm.

Preservation Method: Filtered through both Whatman glass fiber filter (.7µm pore size) and MF-Millipore (.45µm pore size) membrane filter, and stored at or below -20°C.

Instrumentation: AA3 Autoanalyzer made my SEAL Analytical.

c) **Parameters: NO23F**

Method Reference: A modification of the Armstrong et al. procedure is used for the analysis of nitrate and nitrite. Armstrong, F.A.J., Stearns, C.A., and Strickland, J.D.H., 1967. "The measurement of upwelling and subsequent biological processes by means of the Technicon Autoanalyzer and associated equipment," Deep-Sea Research, 14, pp.381-389.

Method Descriptor: For the nitrate analysis, a water sample is passed through a cadmium column where the nitrate is reduced to nitrite. This nitrite is then diazotized with sulfanilamide and coupled with N-(1-naphthyl)-ethylenediamine to form a red dye. The sample is then passed through a 10mm flowcell and the absorbance is measured at 540nm. The procedure is the same for the nitrite analysis but without the cadmium column.

Preservation Method: Filtered through both Whatman glass fiber filter (.7µm pore size) and MF-Millipore (.45µm pore size) membrane filter, and stored at or below -20°C.

Instrumentation: AA3 Autoanalyzer made my SEAL Analytical.

d) **Parameter: CHLA**

Method Reference: Chlorophyll was analyzed using the California Cooperative Oceanic Fisheries Investigations (CalCOFI) methods, which is based on several references such as:

Holm_Hansen, O., Lorenzen, C.J., Holms, R.W., Strickland, J.D.H. (1965). Fluorometric Determination of Chlorophyll. J. Cons.perm.int Explor. Mer. 30: 3-15.

Lorenzen, C. J. (1967) Determination of chlorophylls and phaeopigments: spectrophotometric equations. *Limnol. Oceanogr.* 12: 343–346.

Yentsch, C.S., Menzel, D.W. (1963). A method for the determination of phytoplankton chlorophyll and phaeophytin by fluorescence. *Deep-Sea Res.* 10: 221-231.

Method Descriptor: Samples are filtered through Whatman .7µm particle retention glass fiber filters. Chlorophyll *a* is extracted in an acetone solution. Chlorophyll and phaeopigments are then measured fluorometrically using an acidification technique.

Preservation Method: Filtered through a Whatman .7µm particle retention glass fiber filter and stored in a container with desiccant at or below -20°C.

Instrumentation: Turner 10-005R fluorometer

e) **Parameter: SiO4F**

Method Reference: Silicate is analyzed using the basic method of Armstrong et al. Armstrong, F.A.J., Stearns, C.A., and Strickland, J.D.H., 1967. "The measurement of upwelling and subsequent biological processes by means of the Technicon Autoanalyzer and associated equipment," *Deep-Sea Research*, 14, pp.381-389.

Method Descriptor: Acidified ammonium molybdate is added to a water sample to produce silicomolybdic acid which is then reduced to silicomolybdous acid (a blue compound) following the addition of stannous chloride. The sample is passed through a 10mm flowcell and measured at 660nm.

Preservation Method: Filtered through both Whatman glass fiber filter (.7µm pore size) and MF-Millipore (.45µm pore size) membrane filter, and stored at or below -20°C.

Instrumentation: AA3 Autoanalyzer made by SEAL Analytical.

14) Field and Laboratory QAQC programs –

a) **Precision**

i) **Field variability** – Duplicate samples (i.e., two successive grab samples in different bottles) were collected at all sites.

ii) **Laboratory variability** – NONE

iii) **Inter-organizational splits** – NONE

b) **Accuracy**

i) **Sample spikes** – NONE

ii) **Standard reference material analysis** – Accuracy is based on the quality of the following standards, supplied from Johnson Matthey Chemical Co. and Fisher Scientific. Primary standards for silicate (Na₂SiF₆), nitrate (KNO₃), nitrite (NaNO₂), phosphate (KH₂PO₄), and ammonia (NH₃) have purities of >99%, 99.999%, 98%, 99.999% and > 99.0%, respectively, as reported by the supplier.

Reference materials for nutrients in seawater (RMNS) are also used as a check sample and run periodically (approximately once a month). The RMNS preparation, verification, and suggested protocol for use of the material are described by (Aoyama et al., 2006, 2007, 2008) and Sato (2010). See the end of this document for these citations.

Results of these analyses are within the stated accuracy for the reference materials.

iii) **Cross calibration exercises** - NONE

15) QAQC flag definitions –

QAQC flags provide documentation of the data and are applied to individual data points by insertion into the parameter's associated flag column (header preceded by an F_). QAQC flags are applied to the nutrient data during secondary QAQC to indicate data that are out of sensor range low (-4), rejected due to QAQC checks (-3), missing (-2), optional and were not collected (-1), suspect (1), and that have been corrected (5). All remaining data are flagged as having passed initial QAQC checks (0) when the data are uploaded and assimilated into the CDMO ODIS as provisional plus data. The historical data flag (4) is used to indicate data that were submitted to the CDMO prior to the initiation of secondary QAQC flags and codes (and the use of the automated primary QAQC system for WQ and MET data). This flag is only present in historical data that are exported from the CDMO ODIS.

- 4 Outside Low Sensor Range
- 3 Data Rejected due to QAQC
- 2 Missing Data
- 1 Optional SWMP Supported Parameter
- 0 Data Passed Initial QAQC Checks
- 1 Suspect Data
- 4 Historical Data: Pre-Auto QAQC
- 5 Corrected Data

16) QAQC code definitions –

QAQC codes are used in conjunction with QAQC flags to provide further documentation of the data and are also applied by insertion into the associated flag column. There are three (3) different code categories, general, sensor, and comment. General errors document general problems with the sample or sample collection, sensor errors document common sensor or parameter specific problems, and comment codes are used to further document conditions or a problem with the data. Only one general or sensor error and one comment code can be applied to a particular data point. However, a record flag column (F_Record) in the nutrient data allows multiple comment codes to be applied to the entire data record.

General errors

- | | |
|-----|---|
| GCM | Calculated value could not be determined due to missing data |
| GCR | Calculated value could not be determined due to rejected data |
| GDM | Data missing or sample never collected |
| GQD | Data rejected due to QA/QC checks |
| GQS | Data suspect due to QA/QC checks |
| GSM | See metadata |

Sensor errors

- | | |
|-----|---|
| SBL | Value below minimum limit of method detection |
| SCB | Calculated value could not be determined due to a below MDL component |
| SCC | Calculation with this component resulted in a negative value |
| SNV | Calculated value is negative |

SRD	Replicate values differ substantially
SUL	Value above upper limit of method detection

Parameter Comments

CAB	Algal bloom
CDR	Sample diluted and rerun
CHB	Sample held beyond specified holding time
CIP	Ice present in sample vicinity
CIF	Flotsam present in sample vicinity
CLE	Sample collected later/earlier than scheduled
CRE	Significant rain event
CSM	See metadata
CUS	Lab analysis from unpreserved sample

Record comments

CAB	Algal bloom
CHB	Sample held beyond specified holding time
CIP	Ice present in sample vicinity
CIF	Flotsam present in sample vicinity
CLE	Sample collected later/earlier than scheduled
CRE	Significant rain event
CSM	See metadata
CUS	Lab analysis from unpreserved sample

Cloud cover

CCL	clear (0-10%)
CSP	scattered to partly cloudy (10-50%)
CPB	partly to broken (50-90%)
COC	overcast (>90%)
CFY	foggy
CHY	hazy
CCC	cloud (no percentage)

Precipitation

PNP	none
PDR	drizzle
PLR	light rain
PHR	heavy rain
PSQ	squally
PFQ	frozen precipitation (sleet/snow/freezing rain)
PSR	mixed rain and snow

Tide stage

TSE	ebb tide
TSF	flood tide
TSH	high tide
TSL	low tide

Wave height

WH0	0 to <0.1 meters
WH1	0.1 to 0.3 meters
WH2	0.3 to 0.6 meters
WH3	0.6 to > 1.0 meters
WH4	1.0 to 1.3 meters
WH5	1.3 or greater meters

Wind direction

N	from the north
NNE	from the north northeast

NE	from the northeast
ENE	from the east northeast
E	from the east
ESE	from the east southeast
SE	from the southeast
SSE	from the south southeast
S	from the south
SSW	from the south southwest
SW	from the southwest
WSW	from the west southwest
W	from the west
WNW	from the west northwest
NW	from the northwest
NNW	from the north northwest
<i>Wind speed</i>	
WS0	0 to 1 knot
WS1	> 1 to 10 knots
WS2	> 10 to 20 knots
WS3	> 20 to 30 knots
WS4	> 30 to 40 knots
WS5	> 40 knots

17) Other remarks/notes –

Data may be missing due to problems with sample collection or processing. Laboratories in the NERR System submit data that are censored at a lower detection rate limit, called the Method Detection Limit or MDL. MDLs for specific parameters are listed in the Laboratory Methods and Detection Limits Section (Section II, Part 12) of this document. Concentrations that are less than this limit are censored with the use of a QAQC flag and code, and the reported value is the method detection limit itself rather than a measured value. For example, if the measured concentration of NO₂₃F was 0.0005 mg/l as N (MDL=0.0008), the reported value would be 0.0008 and would be flagged as out of sensor range low (-4) and coded SBL. In addition, if any of the components used to calculate a variable are below the MDL, the calculated variable is removed and flagged/coded -4 SCB. If a calculated value is negative, it is rejected and all measured components are marked suspect. If additional information on MDL's or missing, suspect, or rejected data is needed, contact the Research Coordinator at the reserve submitting the data.

Note: The way below MDL values are handled in the NERRS SWMP dataset was changed in November of 2011. Previously, below MDL data from 2007-2010 were also flagged/coded, but either reported as the measured value or a blank cell. Any 2007-2011 nutrient/pigment data downloaded from the CDMO prior to November of 2011 will reflect this difference.

Sample hold times for 2024: Samples are held at -20°C. NERRS SOP allows nutrient samples to be held for up to 28 days (CHLA for 30) at -20°C, plus allows for up to 5 days for collecting, processing, and shipping samples. Samples held beyond that time period are flagged suspect <1> and coded (CHB). If measured values were below MDL, this resulted in <-4> [SBL] (CHB) flagging/coding.

Sample Descriptor	Date Analyzed				
	PO4F	NH4F	NO23F	CHLA_N	SiO4F
1/23 - 1/24/2024, all diels	2/7/2024	2/7/2024	2/7/2024	2/13/2024	2/7/2024
1/24/2024 all grabs	2/7/2024	2/7/2024	2/7/2024	2/13/2024	2/7/2024
2/20 - 2/21/2024, all diels	3/6/2024	3/6/2024	3/6/2024	3/13/2024	3/6/2024
2/21 - 2/22/2024 all grabs	3/6/2024	3/6/2024	3/6/2024	3/13/2024	3/6/2024
3/18 - 3/19/2024, all diels	4/10/2024	4/10/2024	4/10/2024	4/12/2024	4/10/2024
3/19/2024, all grabs	4/10/2024	4/10/2024	4/10/2024	4/12/2024	4/10/2024
4/16 - 4/17/2024, all diels	5/8/2024	5/8/2024	5/8/2024	5/8/2024	5/8/2024
4/17/2024, all grabs	5/8/2024	5/8/2024	5/8/2024	5/8/2024	5/8/2024
5/15 - 5/16/2024, all diels	5/22/2024	5/22/2024	5/22/2024	5/22/2024	5/22/2024
5/17/2024, all grabs	5/22/2024	5/22/2024	5/22/2024	5/22/2024	5/22/2024
6/18 - 6/19/2024, all diels	6/26/2024	6/26/2024	6/26/2024	6/26/2024	6/26/2024
6/19/2024, all grabs	6/26/2024	6/26/2024	6/26/2024	6/26/2024	6/26/2024
7/22 - 7/23/2024, all diels	8/7/2024	8/7/2024	8/7/2024	7/31/2024	8/7/2024
7/22 - 7/23/2024, all grabs	8/7/2024	8/7/2024	8/7/2024	7/31/2024	8/7/2024
8/19 - 8/20/2024, all diels	8/27/2024	8/27/2024	8/27/2024	8/29/2024	8/27/2024
8/20/2024, all grabs	8/27/2024	8/27/2024	8/27/2024	8/29/2024	8/27/2024
9/16 - 9/17/2024, all diels	9/20/2024	9/20/2024	9/20/2024	10/2/2024	9/20/2024
9/17/2024, all grabs	9/20/2024	9/20/2024	9/20/2024	10/2/2024	9/20/2024
10/15 - 10/16/2024, all diels	10/24/2024	10/24/2024	10/24/2024	10/23/2024	10/24/2024
10/15 - 10/16/2024, all grabs	10/24/2024	10/24/2024	10/24/2024	10/23/2024	10/24/2024
11/12 - 11/13/2024, all diels	11/19/2024	11/19/2024	11/19/2024	11/20/2024	11/19/2024
11/13/2024, all grabs	11/19/2024	11/19/2024	11/19/2024	11/20/2024	11/19/2024
12/10 - 12/11/2024, all diels	1/16/2025*(4 days)	1/16/2025*(4 days)	1/16/2025*(4 days)	12/18/2024	1/16/2025*(4 days)
12/11/2024, all grabs	1/16/2025*(4 days)	1/16/2025*(4 days)	1/16/2025*(4 days)	12/18/2024	1/16/2025*(4 days)

*sample held longer than allowed by NERRS protocols

During storm events, when the flow in the Tijuana River exceeds 1300 liters per second, or during maintenance and/or infrastructure failure, the International Boundary and Water Commission's CILA pump station is shut off. This causes rainwater, urban runoff, and raw sewage to flow over the United States – Mexico border and into the Tijuana River Estuary. As a result, this can disrupt the sample collection process, leading to missed sampling sessions or higher-than-usual nutrient values at the affected sites (Boca Rio and Oneonta Slough). Although we didn't miss any sample sessions this year due to this issue, we had water samples with high nutrient levels throughout 2024. In the case of nutrient-rich samples, they need to be diluted during analysis. Below is a list of dilution ratios and rain events that occurred 72 hours prior a sampling session. Total precipitation values are measured using the tipping bucket from our SWMP Tidal Linkage meteorological station. Data on transboundary flows can be found at the [IBWC Data Portal](#).

Rain events 72 hours prior of sampling:

1/20 – 1/22 1.83in
 2/19 – 2/22 0.43in
 3/15 0.1in
 4/13 0.03in

Specific sample issues:

For the samples below, silica was not reported due to equipment issues

tjrprnut 1/24/2024 13:23
 tjrprnut 1/24/2024 13:24
 tjrsbnut 1/24/2024 12:34
 tjrsbnut 1/24/2024 12:35

Sample dilution details:

tjrbrnut	1/23/2024 12:15	N+N determined by 1:20x dilution; PO4 determined by 1:20x dilution; SIL determined by 1:20x dilution; NO2 determined by 1:20x dilution; NH4 determined by 1:100x dilution
tjrbrnut	1/23/2024 14:30	N+N determined by 1:20x dilution; PO4 determined by 1:20x dilution; SIL determined by 1:20x dilution; NO2 determined by 1:20x dilution; NH4 determined by 1:100x dilution
tjrbrnut	1/23/2024 16:45	N+N determined by 1:20x dilution; PO4 determined by 1:20x dilution; SIL determined by 1:20x dilution; NO2 determined by 1:20x dilution; NH4 determined by 1:100x dilution
tjrbrnut	1/23/2024 19:00	N+N determined by 1:20x dilution; PO4 determined by 1:20x dilution; SIL determined by 1:20x dilution; NO2 determined by 1:20x dilution; NH4 determined by 1:100x dilution
tjrbrnut	1/23/2024 21:15	N+N determined by 1:20x dilution; PO4 determined by 1:20x dilution; SIL determined by 1:20x dilution; NO2 determined by 1:20x dilution; NH4 determined by 1:100x dilution
tjrbrnut	1/23/2024 23:30	N+N determined by 1:30x dilution; PO4 determined by 1:30x dilution; SIL determined by 1:30x dilution; NO2

		determined by 1:30x dilution; NH4 determined by 1:30x dilution
tjrbrnut	1/24/2024 01:45	N+N determined by 1:30x dilution; PO4 determined by 1:30x dilution; SIL determined by 1:30x dilution; NO2 determined by 1:30x dilution; NH4 determined by 1:30x dilution
tjrbrnut	1/24/2024 04:00	N+N determined by 1:30x dilution; PO4 determined by 1:30x dilution; SIL determined by 1:30x dilution; NO2 determined by 1:30x dilution; NH4 determined by 1:100x dilution
tjrbrnut	1/24/2024 06:15	N+N determined by 1:20x dilution; PO4 determined by 1:20x dilution; SIL determined by 1:20x dilution; NO2 determined by 1:20x dilution; NH4 determined by 1:100x dilution
tjrbrnut	1/24/2024 08:30	N+N determined by 1:10x dilution; PO4 determined by 1:10x dilution; SIL determined by 1:10x dilution; NO2 determined by 1:10x dilution; NH4 determined by 1:10x dilution
tjrbrnut	1/24/2024 10:45	N+N determined by 1:20x dilution; PO4 determined by 1:20x dilution; SIL determined by 1:20x dilution; NO2 determined by 1:20x dilution; NH4 determined by 1:20x dilution
tjrbrnut	1/24/2024 13:50	N+N determined by 1:30x dilution; PO4 determined by 1:30x dilution; SIL determined by 1:30x dilution; NO2 determined by 1:30x dilution; NH4 determined by 1:30x dilution
tjrbrnut	1/24/2024 13:51	N+N determined by 1:30x dilution; PO4 determined by 1:30x dilution; SIL determined by 1:30x dilution; NO2 determined by 1:30x dilution; NH4 determined by 1:30x dilution
tjrosnut	1/24/2024 14:33	N+N determined by 1:30x dilution; PO4 determined by 1:30x dilution; SIL determined by 1:30x dilution; NO2 determined by 1:30x dilution; NH4 determined by 1:30x dilution
tjrosnut	1/24/2024 14:34	N+N determined by 1:30x dilution; PO4 determined by 1:30x dilution; SIL determined by 1:30x dilution; NO2 determined by 1:30x dilution; NH4 determined by 1:30x dilution
tjrprnut	1/24/2024 13:23	SIL not reported. Equipment issue - PB, PB2, SB, and SB2 produced some peaks that were questionable.
tjrprnut	1/24/2024 13:24	SIL not reported. Equipment issue - PB, PB2, SB, and SB2 produced some peaks that were questionable.
tjrbsnut	1/24/2024 12:34	SIL not reported. Equipment issue - PB, PB2, SB, and SB2 produced some peaks that were questionable.
tjrbsnut	1/24/2024 12:35	SIL not reported. Equipment issue - PB, PB2, SB, and SB2 produced some peaks that were questionable.
tjrbrnut	2/20/2024 11:30	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrbrnut	2/20/2024 13:45	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution

tjrbrnut	2/20/2024 16:00	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrbrnut	2/20/2024 18:15	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrbrnut	2/20/2024 20:30	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrbrnut	2/20/2024 22:45	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrbrnut	2/21/2024 01:00	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrbrnut	2/21/2024 03:15	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrbrnut	2/21/2024 05:30	NH4 determined by 1:10 dilution
tjrbrnut	2/21/2024 07:45	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrbrnut	2/21/2024 10:00	PO4 determined by 1:30 dilution, NO2 determined by 1:30 dilution, NH4 determined by 1:100 dilution
tjrbrnut	2/22/2024 14:27	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrbrnut	2/22/2024 14:28	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrosnut	2/22/2024 15:00	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrosnut	2/22/2024 15:01	PO4 determined by 1:10 dilution, NO2 determined by 1:10 dilution, NH4 determined by 1:100 dilution
tjrbrnut	3/18/2024 13:00	NO2 determined by 1:6 dilution, NH4 determined by 1:100 dilution
tjrbrnut	3/18/2024 15:15	NH4 determined by 1:100 dilution
tjrbrnut	3/18/2024 17:30	PO4 determined by 1:20 dilution, NH4 determined by 1:100 dilution
tjrbrnut	3/18/2024 19:45	NH4 determined by 1:100 dilution
tjrbrnut	3/18/2024 22:00	NH4 determined by 1:100 dilution
tjrbrnut	3/19/2024 00:15	NH4 determined by 1:100 dilution
tjrbrnut	3/19/2024 02:30	NH4 determined by 1:100 dilution
tjrbrnut	3/19/2024 04:45	NH4 determined by 1:100 dilution
tjrbrnut	3/19/2024 07:00	NH4 determined by 1:100 dilution
tjrbrnut	3/19/2024 09:15	NH4 determined by 1:100 dilution
tjrbrnut	3/19/2024 11:30	NH4 determined by 1:100 dilution
tjrbrnut	3/19/2024 12:33	PO4 determined by 1:20 dilution, NH4 determined by 1:100 dilution
tjrbrnut	3/19/2024 12:34	PO4 determined by 1:20 dilution, NH4 determined by 1:100 dilution
tjrosnut	3/19/2024 14:20	PO4 determined by 1:20 dilution, NH4 determined by 1:100 dilution
tjrosnut	3/19/2024 14:21	PO4 determined by 1:20 dilution, NH4 determined by 1:100 dilution

tjrbrnut	4/16/2024 12:00	PO4 determined by 1:20x dilution, NH4 determined by 1:100x dilution
tjrbrnut	4/16/2024 14:15	PO4 determined by 1:20x dilution, NH4 determined by 1:100x dilution
tjrbrnut	4/16/2024 16:30	PO4 determined by 1:20x dilution, SIL determined by 1:20x dilution, NH4 determined by 1:100x dilution
tjrbrnut	4/16/2024 18:45	NH4 determined by 1:100x dilution
tjrbrnut	4/16/2024 21:00	NH4 determined by 1:100x dilution
tjrbrnut	4/16/2024 23:15	NH4 determined by 1:100x dilution
tjrbrnut	4/17/2024 01:30	NH4 determined by 1:100x dilution
tjrbrnut	4/17/2024 03:45	NH4 determined by 1:100x dilution
tjrbrnut	4/17/2024 06:00	NH4 determined by 1:100x dilution
tjrbrnut	4/17/2024 08:15	NH4 determined by 1:100x dilution
tjrbrnut	4/17/2024 10:30	NH4 determined by 1:100x dilution
tjrbrnut	4/17/2024 10:47	NH4 determined by 1:100x dilution
tjrbrnut	4/17/2024 10:48	NH4 determined by 1:100x dilution
tjrosnut	4/17/2024 10:17	NH4 determined by 1:100x dilution
tjrosnut	4/17/2024 10:18	NH4 determined by 1:100x dilution
tjrbrnut	5/15/2024 11:00	NH4 determined by 1:100x dilution
tjrbrnut	5/15/2024 13:15	NH4 determined by 1:100x dilution
tjrbrnut	5/15/2024 15:30	PO4 determined by 1:100x dilution, NH4 determined by 1:200x dilution, CHLA determined by 1:2 dilution
tjrbrnut	5/15/2024 17:45	NH4 determined by 1:100x dilution
tjrbrnut	5/15/2024 20:00	NH4 determined by 1:100x dilution
tjrbrnut	5/15/2024 22:15	NH4 determined by 1:100x dilution
tjrbrnut	5/16/2024 00:30	NH4 determined by 1:100x dilution
tjrbrnut	5/16/2024 02:45	NH4 determined by 1:100x dilution
tjrbrnut	5/16/2024 05:00	NH4 determined by 1:100x dilution
tjrbrnut	5/16/2024 07:15	NH4 determined by 1:100x dilution
tjrbrnut	5/16/2024 09:30	NH4 determined by 1:100x dilution
tjrbrnut	5/16/2024 11:15	PO4 determined by 1:100x dilution, SIL determined by 1:100x dilution, NH4 determined by 1:200x dilution, CHLA determined by 1:7 dilution
tjrbrnut	5/16/2024 11:16	PO4 determined by 1:100x dilution, SIL determined by 1:100x dilution, NH4 determined by 1:200x dilution, CHLA determined by 1:7 dilution
tjrosnut	5/16/2024 11:50	PO4 determined by 1:100x dilution, SIL determined by 1:100x dilution, NH4 determined by 1:200x dilution, CHLA determined by 1:30 dilution
tjrosnut	5/16/2024 11:51	PO4 determined by 1:100x dilution, SIL determined by 1:100x dilution, NH4 determined by 1:200x dilution, CHLA determined by 1:30 dilution
tjrbrnut	6/18/2024 09:45	PO4 determined by 1:100x dilution, NH4 determined by 1:400x dilution
tjrbrnut	6/18/2024 12:00	PO4 determined by 1:100x dilution, NH4 determined by 1:400x dilution

tjrbrnut	6/18/2024 14:15	PO4 determined by 1:100x dilution, NH4 determined by 1:400x dilution
tjrbrnut	6/18/2024 16:30	NH4 determined by 1:400x dilution
tjrbrnut	6/18/2024 18:45	NH4 determined by 1:100x dilution
tjrbrnut	6/18/2024 21:00	NH4 determined by 1:100x dilution, CHLA determined by 1:8 dilution
tjrbrnut	6/18/2024 23:15	NH4 determined by 1:400x dilution
tjrbrnut	6/19/2024 01:30	PO4 determined by 1:100x dilution, NH4 determined by 1:400x dilution, CHLA determined by 1:8 dilution
tjrbrnut	6/19/2024 03:45	PO4 determined by 1:100x dilution, NH4 determined by 1:400x dilution, CHLA determined by 1:8 dilution
tjrbrnut	6/19/2024 06:00	PO4 determined by 1:100x dilution, NH4 determined by 1:400x dilution, CHLA determined by 1:8 dilution
tjrbrnut	6/19/2024 08:15	NH4 determined by 1:400x dilution
tjrbrnut	6/19/2024 11:01	PO4 determined by 1:100x dilution, NH4 determined by 1:400x dilution
tjrbrnut	6/19/2024 11:02	PO4 determined by 1:100x dilution, NH4 determined by 1:400x dilution
tjrosnut	6/19/2024 11:34	PO4 determined by 1:100x dilution, NH4 determined by 1:400x dilution, CHLA determined by 1:8 dilution
tjrosnut	6/19/2024 11:35	PO4 determined by 1:100x dilution, NH4 determined by 1:400x dilution, CHLA determined by 1:8 dilution
tjrbrnut	7/22/2024 15:15	PO4 determined by 1:10 dilution; NH4 determined by 1:200 dilution
tjrbrnut	7/22/2024 17:30	PO4 determined by 1:10 dilution; NH4 determined by 1:200 dilution
tjrbrnut	7/22/2024 19:45	PO4 determined by 1:10 dilution; NH4 determined by 1:10 dilution
tjrbrnut	7/23/2024 00:15	NH4 determined by 1:100 dilution
tjrbrnut	7/23/2024 02:30	PO4 determined by 1:10 dilution; NH4 determined by 1:10 dilution
tjrbrnut	7/23/2024 04:45	PO4 determined by 1:10 dilution; NH4 determined by 1:30 dilution
tjrbrnut	7/23/2024 09:15	PO4 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrbrnut	7/23/2024 11:30	NH4 determined by 1:30 dilution
tjrbrnut	7/23/2024 13:45	PO4 determined by 1:10 dilution; NH4 determined by 1:10 dilution
tjrbrnut	7/23/2024 14:12	PO4 determined by 1:10 dilution; NH4 determined by 1:200 dilution
tjrbrnut	7/23/2024 14:13	PO4 determined by 1:10 dilution; NH4 determined by 1:200 dilution
tjrosnut	7/23/2024 13:40	PO4 determined by 1:10 dilution; NH4 determined by 1:200 dilution
tjrosnut	7/23/2024 13:41	PO4 determined by 1:10 dilution; NH4 determined by 1:200 dilution
tjrbrnut	8/19/2024 12:15	NO2 & PO4 determined by 1:30 dilution; NH4 determined by 1:100 dilution

tjrbrnut	8/19/2024 14:30	PO4 determined by 1:30 dilution; NH4 determined by 1:200 dilution, CHLA determined by 1:8 dilution
tjrbrnut	8/19/2024 16:45	PO4 determined by 1:30 dilution; NH4 determined by 1:30 dilution
tjrbrnut	8/19/2024 19:00	NH4 determined by 1:30 dilution
tjrbrnut	8/19/2024 21:15	NH4 determined by 1:30 dilution
tjrbrnut	8/19/2024 23:30	PO4 determined by 1:30 dilution; NH4 determined by 1:30 dilution, CHLA determined by 1:7 dilution
tjrbrnut	8/20/2024 01:45	PO4 determined by 1:30 dilution; NH4 determined by 1:30 dilution
tjrbrnut	8/20/2024 04:00	PO4 determined by 1:30 dilution; NH4 determined by 1:100 dilution
tjrbrnut	8/20/2024 06:15	NH4 determined by 1:100 dilution
tjrbrnut	8/20/2024 08:30	NH4 determined by 1:30 dilution
tjrbrnut	8/20/2024 10:45	PO4 determined by 1:30 dilution; NH4 determined by 1:30 dilution
tjrbrnut	8/19/2024 13:57	PO4 determined by 1:30 dilution; NH4 determined by 1:100 dilution
tjrbrnut	8/19/2024 13:58	PO4 determined by 1:30 dilution; NH4 determined by 1:100 dilution
tjrosnut	8/19/2024 13:00	PO4 determined by 1:30 dilution; NH4 determined by 1:100 dilution
tjrosnut	8/19/2024 13:01	PO4 determined by 1:30 dilution; NH4 determined by 1:100 dilution
tjrbrnut	9/16/2024 13:30	PO4 determined by 1:10 dilution; NO2 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrbrnut	9/16/2024 15:45	PO4 determined by 1:10 dilution; NO2 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrbrnut	9/16/2024 18:00	PO4 determined by 1:6 dilution; NH4 determined by 1:6 dilution
tjrbrnut	9/17/2024 00:45	PO4 determined by 1:6 dilution; NO2 determined by 1:6 dilution; NH4 determined by 1:6 dilution
tjrbrnut	9/17/2024 03:00	PO4 determined by 1:10 dilution; NO2 determined by 1:10 dilution; NH4 determined by 1:6 dilution
tjrbrnut	9/17/2024 05:15	PO4 determined by 1:10 dilution; NO2 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrbrnut	9/17/2024 07:30	NH4 determined by 1:10 dilution
tjrbrnut	9/17/2024 09:45	
tjrbrnut	9/17/2024 12:00	PO4 determined by 1:6 dilution; NH4 determined by 1:6 dilution
tjrbrnut	9/16/2024 13:27	PO4 determined by 1:6 dilution; NO2 determined by 1:6 dilution; NH4 determined by 1:6 dilution
tjrbrnut	9/16/2024 13:28	PO4 determined by 1:6 dilution; NO2 determined by 1:6 dilution; NH4 determined by 1:6 dilution
tjrosnut	9/17/2024 11:57	PO4 determined by 1:10 dilution; NO2 determined by 1:10 dilution; NH4 determined by 1:100 dilution

tjrosnut	9/17/2024 11:58	PO4 determined by 1:10 dilution; NO2 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrbrnut	10/15/2024 13:15	NO2, PO4 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrbrnut	10/15/2024 15:30	NO2, PO4 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrbrnut	10/15/2024 17:45	PO4, NH4 determined by 1:10 dilution
tjrbrnut	10/15/2024 20:00	NH4 determined by 1:10 dilution
tjrbrnut	10/16/2024 00:30	NO2, PO4, NH4 determined by 1:10 dilution
tjrbrnut	10/16/2024 02:45	NO2, PO4 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrbrnut	10/16/2024 05:00	NO2, PO4 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrbrnut	10/16/2024 07:15	PO4, NH4 determined by 1:10 dilution
tjrbrnut	10/16/2024 11:45	PO4, NH4 determined by 1:10 dilution
tjrbrnut	10/16/2024 12:00	NO2, PO4 , NH4determined by 1:10 dilution
tjrbrnut	10/16/2024 12:01	PO4, NH4 determined by 1:10 dilution
tjrosnut	10/16/2024 13:00	NO2, PO4 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrosnut	10/16/2024 13:01	NO2, PO4 determined by 1:10 dilution; NH4 determined by 1:100 dilution
tjrbrnut	11/12/2024 12:30	NO2, PO4 & NH4 determined by 1:20 dilution
tjrbrnut	11/12/2024 14:45	NO2, PO4 & NH4 determined by 1:20 dilution
tjrbrnut	11/12/2024 17:00	PO4 determined by 1:20 dilution; NH4 determined by 1:100 dilution
tjrbrnut	11/12/2024 19:15	NH4 determined by 1:30 dilution
tjrbrnut	11/12/2024 21:30	NH4 determined by 1:30 dilution
tjrbrnut	11/12/2024 23:45	NO2, PO4 & NH4 determined by 1:20 dilution
tjrbrnut	11/13/2024 02:00	NO2, PO4 & NH4 determined by 1:20 dilution
tjrbrnut	11/13/2024 04:15	PO4 determined by 1:20 dilution; NH4 determined by 1:100 dilution
tjrbrnut	11/13/2024 06:30	NH4 determined by 1:30 dilution
tjrbrnut	11/13/2024 11:00	NH4 determined by 1:30 dilution
tjrbrnut	11/13/2024 13:30	NO2, PO4 & NH4 determined by 1:20 dilution
tjrbrnut	11/13/2024 13:31	NO2, PO4 & NH4 determined by 1:20 dilution
tjrosnut	11/13/2024 14:00	NO2 & PO4 determined by 1:20 dilution; NH4 determined by 1:100 dilution
tjrosnut	11/13/2024 14:01	NO2 & PO4 determined by 1:20 dilution; NH4 determined by 1:100 dilution
tjrbrnut	12/10/2024 11:30	PO4 determined by 1:10x dilution. NO2 determined by 1:10x dilution. NH4 determined by 1:100x dilution.
tjrbrnut	12/10/2024 13:45	PO4 determined by 1:10x dilution. NO2 determined by 1:10x dilution. NH4 determined by 1:100x dilution.
tjrbrnut	12/10/2024 16:00	NH4 determined by 1:10x dilution.
tjrbrnut	12/10/2024 18:15	NH4 determined by 1:10x dilution.
tjrbrnut	12/10/2024 20:30	PO4 determined by 1:10x dilution. NH4 determined by 1:30x dilution.

tjrbrnut	12/10/2024 22:45	PO4 determined by 1:10x dilution. NH4 determined by 1:30x dilution.
tjrbrnut	12/11/2024 01:00	PO4 determined by 1:10x dilution. NH4 determined by 1:30x dilution.
tjrbrnut	12/11/2024 03:15	NH4 determined by 1:10x dilution.
tjrbrnut	12/11/2024 07:45	NH4 determined by 1:10x dilution.
tjrbrnut	12/11/2024 10:00	PO4 determined by 1:10x dilution. NO2 determined by 1:10x dilution. NH4 determined by 1:30x dilution.
tjrbrnut	12/11/2024 11:53	PO4 determined by 1:10x dilution. NO2 determined by 1:10x dilution. NH4 determined by 1:100x dilution.
tjrbrnut	12/11/2024 11:54	PO4 determined by 1:10x dilution. NO2 determined by 1:10x dilution. NH4 determined by 1:100x dilution.
tjrosnut	12/11/2024 11:28	PO4 determined by 1:10x dilution. NO2 determined by 1:10x dilution. NH4 determined by 1:100x dilution.
tjrosnut	12/11/2024 11:29	PO4 determined by 1:10x dilution. NO2 determined by 1:10x dilution. NH4 determined by 1:100x dilution.